

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International BureauApplicant
copy

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶: A61K 38/06	A1	(11) International Publication Number: WO 96/21456 (43) International Publication Date: 18 July 1996 (18.07.96)
(21) International Application Number: PCT/US96/01559 (22) International Filing Date: 11 January 1996 (11.01.96) (30) Priority Data: 08/371,682 12 January 1995 (12.01.95) US 08/451,839 30 May 1995 (30.05.95) US 08/552,554 3 November 1995 (03.11.95) US 08/582,076 2 January 1996 (02.01.96) US (71) Applicant: UNIVERSITY OF PITTSBURGH [US/US]; 911 William Pitt Union, Pittsburgh, PA 15260 (US). (72) Inventors: SEBTI, Said; 127 Dorseyville Road, Pittsburg, PA 15215 (US). HAMILTON, Andrew; 108 Meadow Heights Drive, Pittsburgh, PA 15215 (US). (74) Agents: KOKULIS, Paul, N. et al.; Cushman, Darby & Cushman, L.L.P., 1100 New York Avenue, N.W., Washington, DC 20005 (US).		(81) Designated States: AU, CA, JP, KR, MX, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: INHIBITORS OF PRENYL TRANSFERASES (57) Abstract Compounds which inhibit prenyl transferases, particularly farnesyltransferase and geranylgeranyl transferase I, processes for preparing the compounds, pharmaceutical compositions containing the compounds, and methods of use.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

INHIBITORS OF PRENYL TRANSFERASES

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to novel
5 peptidomimetics and other compounds which are
useful as inhibitors of protein isoprenyl
transferases (particularly protein
farnesyltransferase and geranylgeranyltransferase)
and as anti-cancer drugs, to compositions
10 containing such compounds and to methods of use.

2. Background Information

Ras proteins are plasma membrane- associated
GTPases that function as relay switches that
transduce biological information from
15 extracellular signals to the nucleus (29-31). In
normal cells Ras proteins cycle between the GDP-
(inactive) and GTP-(active) bound forms to
regulate proliferation and differentiation. The
mechanism by which extracellular signals, such as
20 epidermal and platelet derived growth factor (EGF
and PDGF), transduce their biological information
to the nucleus via Ras proteins has recently been
unraveled (29-31). Binding of the growth factors
to tyrosine kinase receptors results in
25 autophosphorylation of various tyrosines which
then bind src-homology 2 (SH2) domains of several
signaling proteins. One of these, a cytosolic
complex of GRB-2 and a ras exchanger (m-SOS-1), is
recruited by the tyrosine phosphorylated receptor
30 where mSOS-1 catalyzes the exchange of GDP for GTP
on Ras, hence activating it. GTP-bound Ras
recruits Raf, a serine/threonine kinase, to the
plasma membrane where it is activated. Raf
triggers a kinase cascade by phosphorylating
35 mitogen-activated protein (MAP)

kinase/extracellular-regulated protein kinase
(ERK) kinase (MEK) which in turn phosphorylates
MAP Kinase on threonine and tyrosine residues.
Activated MAP Kinase translocates to the nucleus
5 where it phosphorylates transcription factors
(31). Termination of this growth signal is
accomplished by hydrolysis of Ras-GTP to Ras-GDP.

Ras oncogenes are the most frequently
identified activated oncogenes in human tumors (1-
10 3). In a large number of human cancers, Ras is
GTP-locked because of mutations in amino acids 12,
13, or 61 and the above Ras pathway no longer
requires an upstream growth signal and is
uninterrupted. As a consequence, enzymes in this
15 pathway such as Raf, MEK and MAP Kinase are
constitutively activated.

In addition to its inability to hydrolyze
GTP, oncogenic Ras must be plasma membrane-bound
to cause malignant transformation (13). Ras is
20 posttranslationally modified by a lipid group,
farnesyl, which mediates its association with the
plasma membrane (10-14).

Post-translational events leading to membrane
association of p21ras have previously been
25 disclosed (10-14). The p21ras proteins are first
made as pro-p21ras in the cytosol where they are
modified on cysteine 186 of their carboxyl
terminal sequence CA₁A₂X (C = cysteine, A₁ and A₂ =
isoleucine, leucine or valine and X = methionine
30 or serine) by the cholesterol biosynthesis
intermediate farnesyl pyrophosphate (FPP). This
farnesylation reaction is then followed by
peptidase removal of the A₁A₂X tripeptide and
carboxymethylation of the remaining cysteine. The
35 processed p21ras proteins associate with the inner

surface of the plasma membrane (10-14).

5 p21Ras farnesyltransferase, the enzyme responsible for catalyzing the transfer of farnesyl, a 15-carbon isoprenoid, from FPP to the cysteine of the CA₁A₂X carboxyl terminus of p21ras, has been purified to homogeneity from rat brain (15,16). The enzyme is a heterodimer composed of α and β subunits of molecular weights 49 and 46 kDa, respectively (17). The β subunit has been shown to bind p21ras (17). Because p21ras farnesylation and subsequent membrane association are required for p21ras transforming activity (13), it has been proposed that p21ras farnesyltransferase would be a useful anticancer therapy target. Accordingly, an intensive search for inhibitors of the enzyme is underway (18-24, 33-44). Potential inhibitor candidates are CA₁A₂X tetrapeptides which have been shown to be farnesylated by p21ras farnesyltransferase and appear to be potent inhibitors of this enzyme in vitro (15,18,21-24). Competition studies have demonstrated that CA₁A₂X peptides with the greatest inhibitory activity are those where A₁ and A₂ are hydrophobic peptides with charged or hydrophilic residues in the central positions demonstrating very little inhibitory activity (18,21,23). A major drawback with the use of peptides as therapeutic agents is their low cellular uptake and their rapid inactivation by proteases.

30 The research efforts directed towards farnesyltransferase and the inhibition of its activity are further illustrated by the following patents or published patent applications:

U.S. 5,141,851

WO 91/16340

WO 92/18465

EPA 0456180 A1

5


EPA 0461869 A2

EPA 0512865 A2

EPA 0520823 A2

EPA 0523873 A1

10 Of the above disclosures, EPA 0520823 A2 discloses compounds which are useful in the inhibition of farnesyl-protein transferase and the farnesylation of the oncogene protein ras. The compounds of EPA 0520823 A2 are illustrated by the formula:




15 or pharmaceutically acceptable salts thereof, wherein Cys is a cysteine amino acid; Xaa¹ is an amino acid in natural L-isomer form; dXaa² is an amino acid in unnatural D-isomer form; and
20 Xaa³ is an amino acid in natural L-isomer form.

The preferred compounds are said to be CV(Dl)S and CV(Df)M, the amino acids being identified by conventional 3 letter and single letter abbreviations as follows:

25	Cysteine	Cys	C
	Glycine	Gly	G
	Isoleucine	Ile	I
	Leucine	Leu	L
	Lysine	Lys	K
30	Methionine	Met	M
	Phenylalanine	Phe	F
	Serine	Ser	S
	Threonine	Thr	T
	Valine	Val	V

EPA 0523873 A1 discloses a modification of the compounds of EPA 0520823 A2 wherein Xaa³ is phenylalanine or p-fluorophenylalanine.

5 EPA 0461869 describes compounds which inhibit farnesylation of Ras protein of the formula:



where Aaa¹ and Aaa² are aliphatic amino acids and Xaa is an amino acid. The aliphatic amino acids which are disclosed are Ala, Val, Leu and Ile.

10 Preferred compounds are those where Aaa¹ is Val, Aaa² is Leu, Ile or Val and Xaa is Ser or Met. Preferred specific compounds are:

Cys-Val-Leu-Ser

Cys-Val-Ile-Met

15 Cys-Val-Val-Met

U.S. patent 5,141,851 and WO 91/16340 disclose the purified farnesyl protein transferase and certain peptide inhibitors therefor, including, for example, CVIM, TKCVIM and
20 KKSKTKCVIM.

WO 92/18465 discloses certain farnesyl compounds which inhibit the enzymatic methylation of proteins including ras proteins.

EPA 0456180 A1 is directed to a
25 farnesylprotein transferase assay which can be used to identify substances that block farnesylation of ras oncogene gene products while EPA 0512865 A2 discloses certain cyclic compounds that are useful for lowering cholesterol and
30 inhibiting farnesylprotein transferase.

As will be evident from the foregoing, there is a great deal of research effort directed towards the development of inhibitors of farnesyltransferase. However, there still remains

a need for improvements in this critically important area.

An enzyme closely related to farnesyltransferase, geranylgeranyltransferase I (GGTase I), attaches the lipid geranylgeranyl to the cysteine of the CAAX box of proteins where X is leucine (49,69). FTase and GGTase I are α/β heterodimers that share the α subunit (61,62). Cross-linking experiments suggested that both substrates (FPP and Ras CAAX) interact with the β subunit of FTase (17,63). Although GGTase I prefers leucine at the X position, its substrate specificity was shown to overlap with that of FTase in vitro (64). Furthermore, GGTase I is also able to transfer farnesyl to a leucine terminating peptide (65).

Although CAAX peptides are potent competitive inhibitors of FTase, rapid degradation and low cellular uptake limit their use as therapeutic agents. The strategy of the present invention to develop superior compounds for inhibiting FTase and GGTase has been to replace several amino acids in the CAAX motif by peptidemimics. The rationale behind this strategy is based on the existence of a hydrophobic pocket at the enzyme active site that interacts with the hydrophobic "AA" dipeptide of the carboxyl terminus CAAX of Ras molecules. In this regard, two very potent inhibitors of FTase (i.e. Cys-3AMBA-Met and Cys-4ABA-Met) were disclosed by us in an earlier U.S. patent application. The peptidomimetic Cys-4ABA-Met incorporates a hydrophobic/aromatic spacer (i.e. 4-aminobenzoic acid) between Cys and Met. The present application discloses several derivatives of Cys-4ABA-Met where positions 2 and 3 of 4-amino





benzoic acid were modified by several alkyl, and/or aromatic groups, compounds that show great promise of ability to selectively antagonize RAS-dependent signaling and to selectively inhibit the growth of human tumors with aberrant Ras function.

Of the four types of Ras proteins (H-, N-, K4A-, and K4B-Ras) expressed by mammalian cells, K4B-Ras (also called K-Ras4B) is the most frequently mutated form of Ras in human cancers (1,3). Although several laboratories have demonstrated potent inhibition of oncogenic H-Ras processing and signaling (43,44), this disruption has not been shown with K-Ras4B. Previous studies have targeted H-Ras and not K-Ras4B as a target for the development of inhibitors. One recent report indicates that K-Ras4B can be geranylgeranylated *in vitro*, but with relatively low efficiency; its K_m for GGTase I is 7 times higher than its K_m for FTase (67). GGTase I CAAX-based inhibitors that can block geranylgeranylation processing have not been reported.

Recently, we have shown that a potent inhibitor of FTase disrupts K-Ras4B processing but only at very high concentrations that also inhibited the processing of geranylgeranylated proteins (66). This suggested that K-Ras4B may be geranylgeranylated, and that therefore inhibitors targeted at GGTase I would be effective in disrupting oncogenic K-Ras4B processing and signalling, and in treatment of cancers which were related to this form of Ras.

Summary of the Invention

In accordance with the present invention there are compounds of the formula (A-L):




wherein R_1' is

- i) hydrogen;
- ii) lower alkyl;
- iii) alkenyl;
- 5 iv) alkoxy;
- v) thioalkoxy;
- vi) halo;
- vii) haloalkyl;
- viii) aryl- L_2 -, wherein L_2 is absent, $-CH_2-$, $-CH_2CH_2-$, $-CH(CH_3)-$, $-O-$, $-S(O)_q$ wherein q is 0, 1, or 2, $-N(R')$ - wherein R' is hydrogen or lower alkyl, or $-C(O)-$ and aryl is selected from the group consisting of phenyl, naphthyl, tetrahydronaphthyl, indanyl and indenyl and the
- 10 aryl group is unsubstituted or substituted; or
- ix) heterocyclic- L_3 - wherein L_3 is absent, $-CH_2-$, $-CH_2CH_2-$, $-CH(CH_3)-$, $-O-$, $-S(O)_q$ wherein q is 0, 1 or 2, $-N(R')$ - wherein R' is hydrogen or loweralkyl, or $-C(O)-$ and heterocyclic is a
- 15 monocyclic heterocyclic wherein the heterocyclic is unsubstituted or substituted with one, two, or three substituents independently selected from the group consisting of loweralkyl, hydroxy, hydroxyalkyl, halo, nitro, oxo ($=O$), amino, N-protected amino, alkoxy, thioalkoxy and haloalkyl;
- 20
- 25

R_{1a} is hydrogen or lower alkyl;

R_2' is

i)




- 30 wherein R_{12a} is hydrogen, loweralkyl or $-C(O)O-R_{13}$, wherein R_{13} is hydrogen or a carboxy-protecting

group and R_{12b} is hydrogen or loweralkyl, with the proviso that R_{12a} and R_{12b} are not both hydrogen,

ii) $-C(O)NH-CH(R_{14})-C(O)OR_{15}$ wherein R_{14} is

- a) loweralkyl,
- 5 b) cycloalkyl,
- c) cycloalkylalkyl,
- d) alkoxyalkyl,
- e) thioalkoxyalkyl,
- f) hydroxyalkyl,
- 10 g) aminoalkyl,
- h) carboxyalkyl,
- i) alkoxycarbonylalkyl,
- j) arylalkyl or
- k) alkylsulfonylalkyl and

15 R_{15} is hydrogen or a carboxy-protecting group or



R_3' is



where

A represents O or 2H, and

25 R_0 represents SH, NH_2 , or $C_xH_y-SO_2-NH-$, wherein C_xH_y is a straight chain saturated or unsaturated hydrocarbon, with x being between 1 and 20 and y between 3 and 41, inclusive; and

L_1 is -NH-;
or pharmaceutically acceptable salts or prodrugs thereof.


5 An important embodiment of the present invention is based on the finding that a novel group of peptidomimetics as represented by Formula (I) have a high inhibitory potency against human tumor p21ras farnesyltransferase and inhibit tumor growth of human carcinomas:

10 $C\beta X$ (I)

where


C stands for the cysteine radical, or for the reduced form of the cysteine radical (R-2-amino-3-mercaptopropyl amine); β is the radical of a non-peptide aminoalkyl- or amino-substituted phenyl
15 carboxylic acid; and X is the radical of an amino acid, preferably Met. Any other natural or synthetic amino acid can also be used at this position. The invention also includes
20 pharmaceutically acceptable salts and prodrugs of Formula (I).

A particularly preferred compound in this regard is:



In this compound the cysteine radical is in the reduced form and the spacer group is 2-phenyl-4-aminobenzoic acid.

5 Another preferred compound of the invention is:



The compounds of Formula (I) are different from the prior art farnesyltransferase inhibitors in that they do not include separate peptide amino acids A₁, A₂ as in prior art inhibitors represented by the formula CA₁A₂X. The present compounds are consequently free from peptidic amide bonds.

15 It is also to be noted that the present compounds are not farnesylated by the enzyme. They are, therefore, true inhibitors, not just alternative substrates. This may explain the high inhibitory action of the present compounds relative to their parent compounds which are farnesylated.

20 A further important feature of the invention is the provision of the compounds of Formula (I) in the form of pro-drugs. Broadly speaking, this is accomplished by functionalizing the terminal end groups (amino, cysteine sulfur and carboxy groups) of the compounds with hydrophobic, enzyme-sensitive moieties which serve to increase the plasma membrane permeability and cellular uptake of the compounds and consequently their efficiency in inhibiting tumor cell growth. In addition,

prodrugs for amino and cysteine sulfur groups can include loweralkylcarbonyl, arylcarbonyl, arylalkylcarbonyl, alkoxycarbonyl, aryloxycarbonyl, cycloalkylcarbonyl, cycloalkoxycarbonyl, and other groups well known to those skilled in the art.

In this regard, a particularly preferred compound of the invention is the methylester form of FTI-276, which is illustrated in Figure 1A. The above-mentioned pro-drug aspect of the invention is applicable not only to the compounds of the invention but also to prior peptide inhibitors CA_1A_2X as well as any other peptide with potential for biological uses for the purpose of improving the overall effectiveness of such compounds, as hereinafter described.

A further modification involves the provision of CA_1A_2X tetrapeptides or $C\beta X$ peptidomimetics which have been modified by functionalizing the sulfhydryl group of the cysteine C with an alkyl phosphonate substituent, as hereinafter described.

Another important embodiment of the invention contemplates replacing the A_1A_2X portion of the CA_1A_2X tetrapeptide inhibitors with a non-amino acid component while retaining the desired farnesyltransferase inhibiting activity. These compounds may be illustrated by Formula (II):

$$CA$$
$$(II)$$

where C is cysteine or reduced cysteine and Δ represents an aryl or heterocyclic substituent such as 3-aminomethyl-biphenyl-3'-carboxylic acid, which does not include a peptide amino acid but corresponds essentially in size with A_1A_2X , as hereinafter described. The invention also includes pharmaceutically acceptable salts and prodrugs of Formula (II).

The invention also includes compounds in which further substitutions have been made at the cysteine position. These compounds comprise free cysteine thiol and/or terminal amino groups at one end and include a carboxylic acid or carboxylate group at the other end, the carboxylic acid or carboxylate group being separated from the cysteine thiol and/or terminal amino group by a hydrophobic spacer moiety which is free from any linking amido group as in prior CAAX mimetics. As with other compounds of the invention, these compounds are not subject to proteolytic degradation inside cells while retaining the structural features required for FTase inhibition. The compounds selectively inhibit FTase both in vitro and in vivo and offer a number of other advantages over prior CAAX peptide mimetics.

Compounds of this embodiment may be illustrated by the formula:




where C^0 is



A represents O or 2H, and

R_0 represents SH, NH_2 , or $C_xH_y-SO_2-NH-$, wherein C_xH_y is a straight chain saturated or unsaturated hydrocarbon, with x being between 1 and 20 and y between 3 and 41, inclusive; and B stands for $-NHR$, where R is an aryl group. The invention also includes pharmaceutically acceptable salts and prodrugs of Formula (III). In one preferred embodiment of the invention, R is a biphenyl substituted with one or more $-COOH$

groups and/or lower alkyl, e.g., methyl, as represented by the formula:



where R_1 and R_3 represent H or COOH; R_2 represents H, COOH, CH_3 , or COOCH_3 ; R_4 represents H or OCH_3 ; and A represents 2H or O. This formula represents a series of 4-amino-3'-carboxybiphenyl derivatives which mimic the Val-Ile-Met tripeptide but have restricted conformational flexibility. Reduction of the cysteine amide bond (where A is H,H) provides a completely non-peptidic Ras CAAX mimetic.

Preferably, R is a biphenyl group with a -COOH substitution in the 3'- or 4'-position, most preferably the 3'-position, with respect to the NH-aryl group. The -COOH substituent may appear as such or in pharmaceutically acceptable salt or ester form, e.g., as the alkali metal salt or methyl ester.

The features of the invention are illustrated herein by reference to the CAAX tetrapeptide known as CVIM (see EP 0461869 and U.S. Patent 5,141,851) and C-4ABA-M. These compounds are, respectively, Cys-Val-Ile-Met and Cys-4 aminobenzoic acid-Met where Cys is the cysteine radical and Met is the methionine radical.

A preferred non-peptide CAAX mimetic of the invention is reduced cys-4-amino-3'-biphenylcarboxylate identified as 4 in Figure 12,

which is also designated FTI-265. This derivative contains no amide bonds and thus is a true non-peptide mimic of the CAAX tetrapeptide.

5 The compounds of the invention may be used in the carboxylic acid form or as pharmaceutically acceptable salts or esters thereof. Lower alkyl esters are preferred although other ester forms, e.g., phenyl esters, may also be used.

10 It is also an object of the present invention to provide a CAAX peptidomimetic that inhibits GGTase I.

Accordingly, it is an object of the present invention to provide a substance and means of disrupting oncogenic K-Ras4B processing and
15 signaling that affects geranylgeranylation and/or farnesylation processing.

It is a further object of the invention to provide a pharmaceutical composition for treating cancer which is responsive to geranylgeranyl
20 transferase inhibitors, such as, but not limited to, pancreatic and colon cancer.

The latter objects are accomplished by replacing the central "AA" of CAAX tetrapeptides by a hydrophobic spacer and incorporating a
25 leucine or isoleucine residue in the C-terminal position to optimize recognition by GGTase I. Additionally, the cysteine moiety may be replaced by reduced cysteine, or by other functional groups as hereinafter disclosed.

30 An important embodiment of the present invention is based on the finding that a novel group of peptidomimetics as represented by Formula (IV) have a high inhibitory potency against geranylgeranyl transferase and disrupt oncogenic
35 K-Ras4B processing and signalling:

C β L

(IV)

where

C stands for the cysteine radical, or for the reduced form of the cysteine radical (R-2-amino-3-mercaptopropyl amine); β is the radical of a non-peptide aminoalkyl- or amino-substituted phenyl carboxylic acid; and L is the radical of leucine or isoleucine. The invention also includes pharmaceutically acceptable salts and prodrugs of the compounds of Formula (IV).

Preferred compounds of this embodiment are derivatives of Cys-4ABA-Leu which are substituted at the 2 and/or 3 positions of the phenyl ring of 4-aminobenzoic acid (4ABA). The substitutions at these positions include, but are not limited to alkyl, alkoxy and aryl (particularly to straight chain or branched groups of 1-10 carbons of the aforementioned) and naphthyl, heterocyclic rings and heteroaromatic rings.

A particularly preferred compound of this aspect of the invention, GGTI-287, is illustrated in Figure 17. In this compound the cysteine radical is in the reduced form and the spacer group is 2-phenyl-4-aminobenzoic acid. Another preferred compound, also shown in Figure 17, is GGTI-297, which contains the spacer group 2-naphthyl-4-aminobenzoic acid. Other spacer groups which will be readily evident as useful are described herein in connection with farnesyl-transferase inhibitors.

A further important feature of the invention is the provision of the compounds of the invention in the form of pro-drugs. By "pro-drug" is meant a compound to which *in vivo* modification occurs to produce the active compound. Such compounds may, for example, be more readily delivered to their sites of action as pro-drugs. Broadly speaking, the pro-drugs of the instant invention are

produced by functionalizing the terminal end groups (amino, cysteine sulfur and carboxy groups) of the compounds with hydrophobic, enzyme-sensitive moieties which serve to increase the plasma membrane permeability and cellular uptake of the compounds and consequently their efficiency in inhibiting tumor cell growth.

In this regard, a particularly preferred compound of the invention is the methylester form of GGTI-287, GGTI-286, also illustrated in Figure 17.

The compounds of the invention may be used in the same manner as prior CAAX tetrapeptide inhibitors to inhibit p21ras farnesyltransferase or geranylgeranyl transferase in any host containing these enzymes. This includes both in vitro and in vivo use. Compounds which inhibit farnesyltransferase, notably human tumor p21ras farnesyltransferase, and consequently inhibit the farnesylation of the oncogene protein Ras, may be used in the treatment of cancer or cancer cells. It is noted that many human cancers have activated ras and, as typical of such cancers, there may be mentioned colorectal carcinoma, myeloid leukemias, exocrine pancreatic carcinoma and the like. Likewise, compounds which inhibit geranylgeranyl transferase may be used in the treatment of cancer which is related to K-Ras4B.

The compounds of the invention may be used in pharmaceutical compositions of conventional form suitable for oral, subcutaneous, intravenous, intraperitoneal or intramuscular administration to a mammal or host. This includes, for example, tablets or capsules, sterile solutions or suspensions comprising one or more compounds of the invention with a pharmaceutically acceptable

carrier and with or without other additives. Typical carriers for tablet or capsule use include, for example, lactose or corn starch. For oral compositions, aqueous suspensions may be used with conventional suspending agents, flavoring agents and the like.

The amount of inhibitor administered to obtain the desired inhibitory effect will vary but can be readily determined. It is expected that the compounds of the present invention will be administered to humans or other mammals as pharmaceutical or chemotherapeutic agents in dosages of .1 to 1000 mg/kg body weight, preferably 1 to 500 mg/kg body weight and most preferably 10-50 mg/kg body weight. The required dose for a given individual or disease will vary, but can be determined by ordinary skilled practitioners using routine methods. The compounds may be administered via methods well known in the pharmaceutical and medical arts, which include, but are not limited to oral, parenteral, topical, and respiratory (inhalation) routes. Pharmaceutical preparations may contain suitable carriers or diluents. Means of determining suitable carriers and diluents are well known in the pharmaceutical arts.

The term "carboxy protecting group", as used herein, refers to a carboxylic acid protecting ester group employed to block or protect the carboxylic acid functionally while the reactions involving other functional sites of the compound are carried out. Carboxy protecting groups are disclosed in Greene, "Protective Groups in Organic Synthesis", pp. 152-186 (1981), which is hereby incorporated herein by reference. In addition, a carboxy protecting group can be used as a prodrug whereby the carboxy protecting group can be

readily cleaved *in vivo*, for example by enzymatic hydrolysis, to release the biologically active parent. A comprehensive discussion of the prodrug concept is provided by T. Higuchi and V. Stella in

5 "Prodrugs as Novel Delivery Systems", vol. 14 of the ACS Symposium Series, American Chemical Society (1975), which is hereby incorporated by reference. Such carboxy protecting groups are

10 will known to those skilled in the art, having been extensively used in the protection of carboxyl groups in the penicillin and cephalosporin fields, as described in U.S. Pat. No. 3,840,556 and 3,719,667, the disclosures of which are hereby incorporated by reference.

15 Examples of esters useful as prodrugs for compounds containing carboxyl groups can be found on pages 14-21 of "Bioreversible Carriers in Drug Design: Theory and Application", edited by E.B. Roche, Pergamon Press, New York, (1987) which is

20 hereby incorporated by reference. Representative carboxy protecting groups are C1 to C8 loweralkyl (e.g. methyl, ethyl or tertiary butyl and the like); arylalkyl, for example, phenethyl or benzyl and substituted derivatives thereof, for example 5-

25 indanyl and the like; dialkylaminoalkyl (e.g. dimethylaminoethyl and the like); alkanoyloxyalkyl groups such as acetoxymethyl, butyryloxymethyl, valeryloxymethyl, isobutyryloxymethyl, isovaleryloxymethyl, 1-(propionyloxy)-1-ethyl, 1-

30 (pivaloyloxyl)-1-ethyl, 1-methyl-1-(propionyloxy)-1-ethyl, pivaloyloxymethyl, propionyloxymethyl and the like; cycloalkanoyloxyalkyl groups such as cyclopropylcarbonyloxymethyl, cyclobutylcarbonyloxymethyl,

35 cyclopentylcarbonyloxymethyl, cyclohexylcarbonyloxymethyl and the like; aroyloxyalkyl, such as benzoyloxymethyl,

benzoyloxyethyl and the like;
arylalkylcarbonyloxyalkyl, such as
benzylcarbonyloxymethyl, 2-benzylcarbonyloxyethyl
and the like; alkoxycarbonylalkyl or
5 cycloalkyloxycarbonylalkyl, such as
methoxycarbonylmethyl,
cyclohexyloxycarbonylmethyl, 1-methoxycarbonyl-1-
ethyl, and the like; alkoxycarbonyloxyalkyl or
cycloalkyloxycarbonylalkyl, such as
10 methoxycarbonyloxymethyl, t-
butyloxycarbonyloxymethyl, 1-ethoxycarbonyloxy-1-
ethyl, 1-cyclohexyloxycarbonyloxy-1-ethyl and the
like; aryloxycarbonyloxyalkyl, such as 2-
(phenoxy carbonyloxy)ethyl, 2-(5-
15 indanyloxycarbonyloxy)ethyl and the like;
alkoxyalkylcarbonyloxyalkyl, such as 2-(1-methoxy-
2-methylpropan-2-oyloxy)ethyl and the like;
arylalkyloxycarbonyloxyalkyl, such as 2-
(benzyloxycarbonyloxy)ethyl and the like;
20 arylalkenyloxycarbonyloxyalkyl, such as 2-(3-
phenylpropen-2-yloxycarbonyloxy)ethyl and the
like; alkoxycarbonylaminoalkyl, such as t-
butyloxycarbonylaminomethyl and the like;
alkylaminocarbonylaminoalkyl, such as
25 methylaminocarbonylaminomethyl and the like;
alkanoylaminoalkyl, such as acetylaminomethyl and
the like; heterocycliccarbonyloxyalkyl, such as 4-
methylpiperazinylcarbonyloxymethyl and the like;
dialkylaminocarbonylalkyl, such as
30 dimethylaminocarbonylmethyl,
diethylaminocarbonylmethyl and the like; (5-
(loweralkyl)-2-oxo-1,3-dioxolen-4-yl)alkyl, such
as (5-t-butyl-2-oxo-1,3-dioxolen-4-yl)methyl and
the like; and (5-phenyl-2-oxo-1,3-dioxolen-4-
35 yl)alkyl, such as (5-phenyl-2-oxo-1,3-dioxolen-4-
yl)methyl and the like.

Preferred carboxy-protected compounds of the invention are compounds wherein the protected carboxy group is a loweralkyl, cycloalkyl or arylalkyl ester, for example, methyl ester, ethyl ester, propyl ester, isopropyl ester, butyl ester, sec-butyl ester, isobutyl ester, amyl ester, isoamyl ester, octyl ester, cyclohexyl ester, phenylethyl ester and the like or an alkanoyloxyalkyl, cycloalkanoyloxyalkyl, aroyloxyalkyl or an arylalkylcarbonyloxyalkyl ester.

The term "N-protecting group" or "N-protected" as used herein refers to those groups intended to protect the N-terminus of an amino acid or peptide or to protect an amino group against undesirable reactions during synthetic procedures. Commonly used N-protecting groups are disclosed in Greene, "Protective Groups in Organic Synthesis," (John Wiley & Sons, New York (1981)), which is hereby incorporated herein by reference.

The term "alkanoyl" as used herein refers to $R_2C(O)-O-$ wherein R_2 is a loweralkyl group.

The term "alkanoylaminoalkyl" as used herein refers to a loweralkyl radical to which is appended $R_{71}-NH-$ wherein R_{71} is an alkanoyl group.

The term "alkanoyloxy" as used herein refers to $R_{29}C(O)-O-$ wherein R_{29} is a loweralkyl group.

The term "alkanoyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended an alkanoyloxy group.

The term "alkenyl" as used herein refers to a straight or branched chain hydrocarbon containing from 2 to 10 carbon atoms and also containing at least one carbon-carbon double bond. Examples of alkenyl include $-CH=CH_2$, $-CH_2CH=CH_2$, $-C(CH_3)=CH_2$, $-CH_2CH=CHCH_3$, and the like.

The term "alkenylene" as used herein refers to a divalent group derived from a straight or branched chain hydrocarbon containing from 2 to 10 carbon atoms and also containing at least one carbon-carbon double bond. Examples of alkenylene include
5 -CH=CH-, -CH₂CH=CH-, -C(CH₃)=CH-, -CH₂CH=CHCH₂-, and the like.

The term "alkoxy" as used herein refers to R₃₀O- wherein R₃₀ is loweralkyl as defined above. Representative examples of alkoxy groups include methoxy, ethoxy, t-butoxy and the like.

The term "alkoxyalkoxy" as used herein refers to R₃₁O-R₃₂O- wherein R₃₁ is loweralkyl as defined above and R₃₂ is an alkylene radical. Representative examples of alkoxyalkoxy groups include methoxymethoxy, ethoxymethoxy, t-butoxymethoxy and the like.

The term "alkoxyalkyl" as used herein refers to an alkoxy group as previously defined appended to an alkyl group as previously defined. Examples of alkoxyalkyl include, but are not limited to, methoxymethyl, methoxyethyl, isopropoxymethyl and the like.

The term "alkoxyalkylcarbonyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended R₆₆-C(O)-O- wherein R₆₆ is an alkoxyalkyl group.

The term "alkoxycarbonyl" as used herein refers to an alkoxy group as previously defined appended to the parent molecular moiety through a carbonyl group. Examples of alkoxycarbonyl include methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl and the like.

The term "alkoxycarbonylalkyl" as used herein refers to an alkoxycarbonyl group as previously defined appended to a loweralkyl radical.

Examples of alkoxycarbonylalkyl include methoxycarbonylmethyl, 2-ethoxycarbonylethyl and the like.

5 The term "alkoxycarbonylaminoalkyl" as used herein refers to a loweralkyl radical to which is appended R_{69} -NH- wherein R_{69} is an alkoxycarbonyl group.

10 The term "alkoxycarbonyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended R_{63} -O- wherein R_{63} is an alkoxycarbonyl group.

15 The term "alkylamino" as used herein refers to R_{35} NH- wherein R_{35} is a loweralkyl group, for example, methylamino, ethylamino, butylamino, and the like.

The term "alkylaminoalkyl" as used herein refers a loweralkyl radical to which is appended an alkylamino group.

20 The term "alkylaminocarbonylaminoalkyl" as used herein refers to a loweralkyl radical to which is appended R_{70} -C(O)-NH- wherein R_{70} is an alkylamino group.

25 The term "alkylene" as used herein refers to a divalent group derived from a straight or branched saturated hydrocarbon having from 1 to 10 carbon atoms by the removal of two hydrogen atoms, for example methylene, 1,2-ethylene, 1,1-ethylene, 1,3-propylene, 2,2-dimethylpropylene, and the like.

30 The term "alkylsulfinyl" as used herein refers to R_{33} S(O)- wherein R_{33} is a loweralkyl group.

35 The term "alkylsulfonyl" as used herein refers to R_{34} S(O)₂- wherein R_{34} is a loweralkyl group.

The term "alkylsulfonylalkyl" as used herein refers to a loweralkyl radical to which is appended an alkylsulfonyl group.

5 The term "alkynyl" as used herein refers to a straight or branched chain hydrocarbon containing from 2 to 10 carbon atoms and also containing at least one carbon-carbon triple bond. Examples of alkynyl include $-C\equiv CH$, $-CH_2C\equiv CH$, $-CH_2C\equiv CCH_3$, and the like.

10 The term "alkynylene" as used herein refers to a divalent group derived from a straight or branched chain hydrocarbon containing from 2 to 10 carbon atoms and also containing at least one carbon-carbon triple bond. Examples of alkynylene include

15 $-C\equiv C-$, $-CH_2C\equiv C-$, $-CH_2C\equiv CCH_2-$, and the like.

The term "amino" as used herein refers to $-NH_2$.

20 The term "aminoalkyl" as used herein refers to a loweralkyl radical to which is appended an amino group.

The term "aryloxyalkyl" as used herein refers to a loweralkyl radical to which is appended an aryloxy group (i.e., $R_{61}-C(O)O-$ wherein R_{61} is an aryl group).

25 The term "aryl" as used herein refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like. Aryl groups (including bicyclic aryl groups) can be unsubstituted or substituted with one, two or three substituents independently selected from loweralkyl, haloalkyl, alkoxy, thioalkoxy, amino, alkylamino, dialkylamino, hydroxy, halo, mercapto, nitro, cyano, carboxaldehyde, carboxy, alkoxycarbonyl, haloalkyl-C(O)-NH-, haloalkenyl-C(O)-NH- and

carboxamide. In addition, substituted aryl groups include tetrafluorophenyl and pentafluorophenyl.

5 The term "arylalkenyl" as used herein refers to an alkenyl radical to which is appended an aryl group.

The term "arylalkenyloxycarbonyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended $R_{66}-O-C(O)-O-$ wherein R_{66} is an arylalkenyl group.

10 The term "arylalkyl" as used herein refers to a loweralkyl radical to which is appended an aryl group. Representative arylalkyl groups include benzyl, phenylethyl, hydroxybenzyl, fluorobenzyl, fluorophenylethyl and the like.

15 The term "arylalkylcarbonyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended an arylalkylcarbonyloxy group (i.e., $R_{62}C(O)O-$ wherein R_{62} is an arylalkyl group).

20 The term "arylalkyloxycarbonyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended $R_{67}-O-C(O)-O-$ wherein R_{67} is an arylalkyl group.

25 The term "aryloxyalkyl" as used herein refers to a loweralkyl radical to which is appended $R_{65}-O-$ wherein R_{65} is an aryl group.

The term "aryloxthioalkoxyalkyl" as used herein refers to a loweralkyl radical to which is appended $R_{75}-S-$ wherein R_{75} is an aryloxyalkyl group.

30 The term "aryloxycarbonylalkyl" as used herein refers to a loweralkyl radical to which is appended $R_{65}-O-C(O)-O-$ wherein R_{65} is an aryl group.

The term "arylsulfonyl" as used herein refers to $R_{36}S(O)_2-$ wherein R_{36} is an aryl group.

35 The term "arylsulfonyloxy" as used herein refers to $R_{37}S(O)_2O-$ wherein R_{37} is an aryl group.

The term "carboxyalkyl" as used herein refers to a loweralkyl radical to which is appended a carboxy (-COOH) group.

5 The term "carboxaldehyde" as used herein refers to the group -C(O)H.

The term "carboxamide" as used herein refers to the group -C(O)NH₂.

10 The term "cyanoalkyl" as used herein refers to a loweralkyl radical to which is appended a cyano (-CN) group.

The term "cycloalkanoylalkyl" as used herein refers to a loweralkyl radical to which is appended a cycloalkanoyl group (i.e., R₆₀-C(O)- wherein R₆₀ is a cycloalkyl group).

15 The term "cycloalkanoyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended a cycloalkanoyloxy group (i.e., R₆₀-C(O)O- wherein R₆₀ is a cycloalkyl group).

20 The term "cycloalkenyl" as used herein refers to an alicyclic group comprising from 3 to 10 carbon atoms and containing a carbon-carbon double bond including, but not limited to, cyclopentenyl, cyclohexenyl and the like.

25 The term "cycloalkyl" as used herein refers to an alicyclic group comprising from 3 to 10 carbon atoms including, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, norbornyl, adamantyl and the like.

30 The term "cycloalkylalkyl" as used herein refers to a loweralkyl radical to which is appended a cycloalkyl group. Representative examples of cycloalkylalkyl include cyclopropylmethyl, cyclohexylmethyl, 2-(cyclopropyl)ethyl, adamantylmethyl and the like.

35 The term "cycloalkyloxycarbonyloxyalkyl" as used herein refers to a loweralkyl radical to

which is appended $R_{64}-O-C(O)-O-$ wherein R_{64} is a cycloalkyl group.

5 The term "dialkoxyalkyl" as used herein refers to a loweralkyl radical to which is appended two alkoxy groups.

The term "dialkylamino" as used herein refers to $R_{38}R_{39}N-$ wherein R_{38} and R_{39} are independently selected from loweralkyl, for example, dimethylamino, diethylamino, methyl propylamino, and the like.

10 The term "dialkylaminoalkyl" as used herein refers to a loweralkyl radical to which is appended a dialkylamino group.

15 The term "dialkylaminocarbonylalkyl" as used herein refers to a loweralkyl radical to which is appended $R_{73}-C(O)-$ wherein R_{73} is a dialkylamino group.

20 The term "dioxoalkyl" as used herein refers to a loweralkyl radical which is substituted with two oxo ($=O$) groups.

The term "dithioalkoxyalkyl" as used herein refers to a loweralkyl radical to which is appended two thioalkoxy groups.

25 The term "halogen" or "halo" as used herein refers to I, Br, Cl or F.

The term "haloalkenyl" as used herein refers to an alkenyl radical, as defined above, bearing at least one halogen substituent.

30 The term "haloalkyl" as used herein refers to a lower alkyl radical, as defined above, bearing at least one halogen substituent, for example, chloromethyl, fluoroethyl or trifluoromethyl and the like.

35 The term "heterocyclic ring" or "heterocyclic" or "heterocycle" as used herein refers to a 5-, 6- or 7-membered ring containing one, two or three heteroatoms independently

selected from the group consisting of nitrogen, oxygen and sulfur or a 5-membered ring containing 4 nitrogen atoms; and includes a 5-, 6- or 7-membered ring containing one, two or three

5 nitrogen atoms; one oxygen atom; one sulfur atom; one nitrogen and one sulfur atom; one nitrogen and one oxygen atom; two oxygen atoms in non-adjacent positions; one oxygen and one sulfur atom in non-adjacent positions; two sulfur atoms in non-

10 adjacent positions; two sulfur atoms in adjacent positions and one nitrogen atom; two adjacent nitrogen atoms and one sulfur atom; two non-adjacent nitrogen atoms and one sulfur atom; two non-adjacent nitrogen atoms and one oxygen atom.

15 The 5-membered ring has 0-2 double bonds and the 6- and 7-membered rings have 0-3 double bonds. The term "heterocyclic" also includes bicyclic, tricyclic and tetracyclic groups in which any of the above heterocyclic rings is fused to one or


20 two rings independently selected from the group consisting of an aryl ring, a cyclohexane ring, a cyclohexene ring, a cyclopentane ring, a cyclopentene ring and another monocyclic heterocyclic ring (for example, indolyl, quinolyl, isoquinolyl, tetrahydroquinolyl, benzofuryl or benzothienyl and the like). Heterocyclics

25 include: pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, piperidinyl, homopiperidinyl, pyrazinyl,

30 piperazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiomorpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl,

35 indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzothiazolyl, benzoxazolyl, furyl, thienyl, thiazolidinyl, isothiazolyl,

triazolyl, tetrazolyl, oxadiazolyl, thiadiazolyl,
 pyrimidyl, tetrahydrofuranyl, dihydrofuranyl,
 tetrahydrothienyl, dihydrothienyl, dihydroindolyl,
 tetrahydroquinolyl, tetrahydroisoquinolyl,
 5 pyranyl, dihydropyranyl, dithiazolyl, benzofuranyl
 and benzothienyl. Heterocyclics also include
 bridged bicyclic groups wherein a monocyclic
 heterocyclic group is bridged by an alkylene
 group, for example,




10

like.

and the

Heterocyclics also include compounds of the
 formula



wherein X* is -CH₂-, -CH₂O- or -O- and Y* is -C(O)-
 15 or -(C(R''))_v- wherein R'' is hydrogen or C₁-C₄-alkyl
 and v is 1, 2 or 3 such as 1,3-benzodioxolyl, 1,4-
 benzodioxanyl and the like.

Heterocyclics can be unsubstituted or
 substituted with one, two or three substituents
 20 independently selected from the group consisting
 of

- a) hydroxy,
- b) -SH,
- c) halo,
- 25 d) oxo (=O),
- e) thioxo (=S),
- f) amino,
- g) -NHOH,
- h) alkylamino,
- 30 i) dialkylamino,

- j) alkoxy,
- k) alkoxyalkoxy.
- l) haloalkyl.
- m) hydroxyalkyl,
- 5 n) alkoxyalkyl,
- o) cycloalkyl,
- p) cycloalkenyl,
- q) alkenyl,
- r) alkynyl,
- 10 s) aryl,
- t) arylalkyl,
- u) -COOH,
- v) -SO₃H,
- w) loweralkyl,
- 15 x) alkoxycarbonyl,
- y) -C(O)NH₂,
- z) -C(S)NH₂,
- aa) -C(=N-OH)NH₂,
- bb) loweralkyl-C(O)-,
- 20 cc) loweralkyl-C(S)-,
- dd) formyl,
- ee) cyano, and
- ff) nitro.

25 The term "(heterocyclic)alkyl" as used herein refers to a heterocyclic group as defined above appended to a loweralkyl radical as defined above. Examples of heterocyclic alkyl include 2-pyridylmethyl, 4-pyridylmethyl, 4-quinolinylmethyl and the like.

30 The term "heterocycliccarbonyloxyalkyl" as used herein refers to a loweralkyl radical to which is appended R₇₂-C(O)-O- wherein R₇₂ is a heterocyclic group.

35 The term "hydroxyalkyl" as used herein refers to a loweralkyl radical to which is appended an hydroxy group.

The term "hydroxythioalkoxy" as used herein refers to $R_{51}S-$ wherein R_{51} is a hydroxyalkyl group.

The term "loweralkyl" as used herein refers to branched or straight chain alkyl groups comprising one to ten carbon atoms, including methyl, ethyl, propyl, isopropyl, n-butyl, neopentyl and the like.

The term "N-protected alkylaminoalkyl" as used herein refers to an alkylaminoalkyl group wherein the nitrogen is N-protected.

The term "oxoalkyloxy" as used herein refers to an alkoxy radical wherein the loweralkyl moiety is substituted with an oxo (=O) group.

The term "spiroalkyl" as used herein refers to an alkylene diradical, both ends of which are bonded to the same carbon atom of the parent group to form a spirocyclic group.

The term "thioalkoxy" as used herein refers to $R_{52}S-$ wherein R_{52} is loweralkyl. Examples of thioalkoxy include, but are not limited to, methylthio, ethylthio and the like.

The term "thioalkoxyalkyl" as used herein refers to a thioalkoxy group as previously defined appended to a loweralkyl group as previously defined. Examples of thioalkoxyalkyl include thiomethoxymethyl, 2-thiomethoxyethyl and the like.

The present invention also relates to processes for preparing the compounds of formula (I)-(XI) and to the synthetic intermediates useful in such processes.

In a further aspect of the present invention are disclosed pharmaceutical compositions which comprise a compound of the present invention in combination with a pharmaceutically acceptable carrier.

In yet another aspect of the present invention are disclosed pharmaceutical compositions which comprise a compound of the present invention in combination with another
5 chemotherapeutic agent and a pharmaceutically acceptable carrier.

In yet another aspect of the present invention is disclosed a method for inhibiting protein isoprenyl transferases (i.e., protein
10 farnesyltransferase and/or geranylgeranyltransferase) in a human or lower mammal, comprising administering to the patient a therapeutically effective amount of a compound of the invention.

15 In yet another aspect of the present invention is disclosed a method for inhibiting post-translational modification of the oncogenic Ras protein by protein farnesyltransferase, protein geranylgeranyltransferase or both.

20 In yet another aspect of the present invention is disclosed a method for treatment of conditions mediated by farnesylated or geranylgeranylated proteins, for example, treatment of Ras associated tumors in humans and
25 other mammals.

In yet another aspect of the present invention is disclosed a method for inhibiting or treating cancer in a human or lower mammal, comprising administering to the patient a
30 therapeutically effective amount of a compound of the invention alone or in combination with another chemotherapeutic agent.

In yet another aspect of the present invention is disclosed a method for treating or
35 preventing restenosis in a human or lower mammal, comprising administering to the patient a

therapeutically effective amount of a compound of the invention.

The compounds of the present invention can be used in the form of pharmaceutically acceptable salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as loweralkyl halides (such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides), dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid.

Basic addition salts can be prepared *in situ* during the final isolation and purification of the

compounds of formulas A-L, or separately by reacting the carboxylic acid function with a suitable base such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia, or an organic primary, secondary or tertiary amine. Such pharmaceutically acceptable salts include, but are not limited to, cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium, aluminum salts and the likes, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. Other representative organic amines useful for the formation of base addition salts include diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like.

Other features of the invention will also be hereinafter apparent.

Brief Description of the Drawings

Figure 1: Ras CAAX peptidomimetics and FTase/GGTase I activities

A. Structures of CVIM, FTI-249, FTI-276 and FTI-277. FTI-276 and FTI-277 were synthesized as described in Examples 10 and 11. B. FTase and GGTase I inhibition assays were carried out as described in Example 12 by determining the ability of FTI-276 to inhibit the transfer of farnesyl and geranylgeranyl to recombinant H-Ras-CVLS and H-Ras-CVLL, respectively. The data are representative of at least three different experiments.

Figure 2: Inhibition of Ras and Rap1A Processing

A. H-RasF cells were treated with various concentrations of FTI-277, lysed and the lysates immunoblotted with anti-Ras or anti-Rap1A antibodies as described in Example 13. B. pZIPneo, H-RasF, H-RasGG, Raf and S186 cells were treated with vehicle or FTI-277 (5 μ M), lysed and lysates immunoblotted by anti-Ras antibody. Data is representative of 5 different experiments. The cells were obtained from Dr. Channing Der, University of North Carolina, Chapel Hill, North Carolina.

Figure 3: Effects of FTI-277 on Ras/Raf

Association. pZIPneo, H-RasF, H-RasGG and S186 cells were treated with vehicle or FTI-277 (5 μ M), homogenized and the membrane (A) and cytosolic (B) fractions were separated and immunoprecipitated by an anti-Raf antibody. The immunoprecipitates were then separated by SDS-PAGE and immunoblotted with anti-RAS antibody as described in Example 14. Data is representative of three different experiments.

Figure 4: Effects of FTI-277 on Ras Nucleotide Binding and Raf Kinase Activity

A: H-RasF cells were treated with vehicle or FTI-277, lysed and the lysates immunoprecipitated with anti-Ras antibody. The GTP and GDP were then released from Ras and separated by TLC as described in Example 15. B: pZIPneo and H-RasF cells were treated with vehicle or FTI-277, lysed and cells lysates immunoprecipitated with an anti-Raf antibody. Raf kinase was assayed by using a 19-mer autophosphorylation peptide as substrate as described in Example 16. Data are representative of three different experiments.

Figure 5: Effect of FTI-277 on Oncogenic Activation of MAPK.

5 A: H-RasF cells were treated with various concentrations of FTI-277, cells lysed and lysates run on SDS-PAGE and immunoblotted with anti-MAPK antibody. B: pZIPneo, H-RasF, H-RasGG, Raf, and S186 cells were treated with vehicle of FTI-277 (5 μ M), lysed and cells lysates processed as for A. Data are representative of two different experiments.

10

Figure 6. FTI-276 inhibits selectively Ras processing and oncogenic Ras activation of MAP Kinase. NIH 3T3 cells transfected with empty vector (pZIPneo), oncogenic (GTP-locked) farnesylated Ras (RasF), geranylgeranylated Ras (RasGG) or a transforming mutant of human Raf-1 were obtained from Channing Der and Adrienne Cox (University of North Carolina, Chapel Hill, NC, USA) (26,27). The cells were plated in DMEM/10% CS (Dubelco's Modified Eagles Medium, 10% calf serum) on day one and treated with vehicle or FTI-270 (20 μ M) on days 2 and 3. The cells were then harvested on day 4 and lysed in lysis buffer (30 mM HEPES, pH 7.5, 1% TX-100, 10% glycerol, 10 mM NaCl, 5 mM MgCl₂, 25 mM NaF, 1 mM EGTA, 2 mM Na₃VO₄, 10 μ g/ml Trypsin inhibitor, 25 μ g/ml leupeptin, 10 μ g/ml aprotinin, 2 mM PMSF). The lysate (35 μ g) was electrophoresed on 15% SDS-PAGE, transferred to nitrocellulose membranes and immunoblotted simultaneously with anti-Ras antibody Y13-238 (isolated from hybridomas purchased from ATCC, Rockville, MD) and an Anti-MAP kinase (erk2) antibody (UBI, Lake Placid, NY) as described previously (17, 22).

15

20

25

30

Figure 7. Antitumor efficacy of FTI-276 against human lung carcinomas. Calu-1 (Panel A) and NCI-H810 cells (Panel B) were purchased from ATCC and grown in McCoy's 5A medium in 10% FBS (Fetal Bovine Serum) and RPMI 1640 in 10% FBS, respectively. The cells were harvested, resuspended in PBS and injected s.c. into the right and left flank of 8 week old female nude mice (10^7 cells/flank). Nude mice (Harlan Sprague Dawley, Indianapolis, Indiana) were maintained in accordance with the Institutional Animal Care and Use Committee (IACUC) procedures and guidelines. On day 32 after s.c. implantation of tumors, animals were dosed i.p. with 0.2 ml once daily for 36 days. Control animals (filled circles) received a saline vehicle whereas treated animals (open triangles) were injected with FTI-276 (50 mg/kg). The tumor volumes were determined by measuring the length (l) and the width (w) and calculating the volume ($V = (l) \times (w)^2/2$). Data are presented as the average volume of eight tumors in each group for each cell line. Statistical significance between control and treated groups were evaluated by using student t test ($*P < 0.05$).

Figure 8. Antitumor Efficacy of FTI-276 and FTI-277 in Human Lung Carcinoma (Calu-1) Cells. Experimental procedure was the same as described in Figure 7.

Figure 9. Inhibition of Tumor Growth in Ras transformed cells by FTI-276 and FTI-277. Ras-transformed NIH 3T3 cells were implanted subcutaneously into nude mice, and daily intraperitoneal injections with FTI-276 and FTI-

277 (50 mg/kg) were started when the tumors reached 50 mm³.

5 **Figure 10. Inhibition of Tumor Growth in Raf transformed cells by FTI-276 and FTI-277.** Raf-transformed NIH 3T3 cells were implanted subcutaneously into nude mice, and daily intraperitoneal injections with FTI-276 and FTI-277 (50 mg/kg) were started when the tumors
10 reached 50 mm³.

Figure 11. Dose response: Antitumor efficacy and Ras processing correlations. A. Antitumor efficacy was carried out as described in Fig. 3 except that animals were randomly assigned to four
15 groups each of 4 mice each (2 tumors per mouse). Saline treated groups (circles); FTI-276 treated groups: 10 mg/kg (squares), 50 mg/kg (upward triangles), 100 mg/kg (downward triangles). B. Ras processing was carried out 5 hours after the
20 last treatment on day 17. Tumors were extracted from the animals, tissued, and lysed in lysis buffer as described in Fig. 1. Lysates (25 µg) were electrophoresed on a 12.5% SDS-PAGE and immunoblotted with anti-Ras antibody Y13-238 as
25 described previously. The blots were then reprobated with anti-Rap1A antibody (Santa Cruze Biotechnologies, Santa Cruz, California).

Figure 12. Structures of CVIM, C-4ABA-M, reduced C-4ABA-M, FTI-265 (4), FTI-271 (5), and FTI-261 (8).
30

Figure 13. Energy-minimized structural conformations for CVIM and farnesyltransferase inhibitor FTI-265.

Figures 14A and B. Comparison of FTase and GGTase I inhibition by FTI-265 and FTI-271.

Figure 15. Silica gel TLC relating to Ras CAAAX peptide and peptidomimetic farnesylation.

5 Figure 16. Ras and Rap1A processing in cells using a compound according to the invention.

Figure 17. CAAAX peptidomimetic structures. Structures of FTI-276/277, GGTI-287/286, and GGTI-297.

10 Figure 18. Disruption of H-Ras and Rap1A processing. NIH 3T3 cells that overexpress oncogenic H-Ras were treated with various concentrations of FTI-277 (0-50 μ M) or GGTI-286 (0-30 μ M). The cells were lysed and the lysates
15 were electrophoresed on SDS-PAGE and immunoblotted with either anti-Ras or anti-Rap1A antibodies as described in Example 3. U and P designate unprocessed and processed forms of the proteins. Data are representative of three independent
20 experiments.

Figure 19. Disruption of K-Ras4B processing. NIH 3T3 cells that overexpress oncogenic K-Ras4B were treated with FTI-277 or GGTI-286 (0-30 μ M). The cells were lysed and the lysates were
25 electrophoresed on SDS-PAGE and immunoblotted with anti-Ras antibodies as described in Example 3. U and P designate unprocessed and processed forms of Ras. The data are representative of three independent experiments.

30 Figure 20. Inhibition of oncogenic activation of MAP Kinase. NIH 3T3 cells that overexpress

either oncogenic H-Ras or K-Ras4B were treated with either FTI-277 or GGTI-286 (0-30 μ M). The cells were lysed and the lysates were electrophoresed on SDS-PAGE and immunoblotted with an anti-MAP kinase antibody. P-MARK designates hyperphosphorylated MAP kinase. The data are representative of three independent experiments.

Figure 21. Inhibition of FTase and GGTase I Activity by GGTI-297.

Figure 22. Antitumor Efficacy of GGTI-286 in K-Ras4B.

Description of Preferred Embodiments

For ease of reference, the following abbreviations may be used in the present specification:

FTase: farnesyltransferase;
GGTase: geranylgeranyltransferase;
SDS-PAGE: sodium dodecyl sulfate
polyacrilamide gel
electrophoresis
PBS: phosphate-buffered saline;
CAAX: tetrapeptide where C is cysteine, A is an aliphatic amino acid and X is an amino acid
DTT: dithiothreitol;
DOC: deoxycholate
BSA: bovine serum albumin
GGTase I: geranylgeranyl transferase I;
PAGE: polyacrylamide gel electrophoresis;
MAPK: mitogen activated protein kinase;
FTI: farnesyltransferase inhibitor;
GGTI: geranylgeranyltransferase inhibitor;

PMSF: phenylmethylsulfonyl fluoride.

I. Farnesyltransferase Inhibitors of the type C β X

The peptidomimetics of Formula (I), one of the preferred embodiments of the invention, may be made using procedures which are conventional in the art. Preferably β is 2-phenyl-4-aminobenzoic acid although constrained derivatives such as tetrahydroisoquinoline-7-carboxylic acid, 2-aminomethyl pyridine-6-carboxylic acid or other heterocyclic derivatives, may also be used. Compounds in which β is an aminomethylbenzoic acid (particularly 3-aminomethylbenzoic acid) are disclosed in U.S. Patent application No. 08/062,287, which is hereby incorporated herein by reference. The acid component of β is conveniently reacted with cysteine so that the amino group of β and the cysteine carboxyl group react to form an amido group, other reactive substituents in the reactants being suitably protected against undesired reaction. In the case of the reduced-cysteine series of compounds, the amino group of β is reacted with a suitably protected cysteinal. The amino acid represented by X, preferably Met, is then reacted through its amino group with the deprotected and activated carboxyl group of spacer compound β . Following deprotection by removal of other protecting groups, the compound of Formula (I) is obtained.

As an alternative, β may first be reacted with the X amino acid followed by reaction with the cysteine or cysteinal component using conventional reaction conditions.

The invention also includes the pharmaceutically acceptable salts of the compounds of Formula (I). These may be obtained by reacting the free base or acid with the appropriate amount

of inorganic or organic acid or base, e.g. an alkali metal hydroxide or carbonate, such as sodium hydroxide, an organic amine, e.g. trimethylamine or the like. Acid salts include the reaction products obtained with, for example, toluene sulfonic acid, acetic acid, propionic acid or the like as conventionally used in the art.

5 The compounds of the invention may be used to inhibit p21ras farnesyltransferase in any host containing the same. This includes both *in vitro* and *in vivo* use. Because the compounds inhibit farnesyltransferase, notably human tumor p21ras farnesyltransferase, and consequently inhibit the farnesylation of the oncogene protein ras, they may be used in the treatment of cancer or cancer cells. It is noted that many human cancers have activated ras and, as typical of such cancers, there may be mentioned colorectal carcinoma, myeloid leukemias, exocrine pancreatic carcinoma and the like.

15 The compounds of the invention may be used in pharmaceutical compositions of conventional form suitable for oral, subcutaneous, intravenous, intraperitoneal or intramuscular administration to a mammal or host. This includes, for example, tablets or capsules, sterile solutions or suspensions comprising one or more compounds of the invention with a pharmaceutically acceptable carrier and with or without other additives. Typical carriers for tablet or capsule use include, for example, lactose or corn starch. For oral compositions, aqueous suspensions may be used with conventional suspending agents, flavoring agents and the like.

30 The amount of inhibitor administered to obtain the desired inhibitory effect will vary but can be readily determined. For human use, daily

dosages are dependent on the circumstances, e.g. age and weight. However, daily dosages of from 0.1 to 20 mg per kg body weight may be mentioned for purposes of illustration.


5 The various aspects of the invention are further described by reference to the following examples. These examples illustrate, among other things, the preparation of the present peptidomimetics and compounds compared therewith.


10 In the invention, the β component is, in general, any non-peptide amino acid combination or other hydrophobic spacer element that produces a compound which mimics the structure and conformation of CVIM or like tetrapeptides CA_1A_2X .
15 A variety of hydrophobic spacers have been used as the β component according to this aspect of the invention. This includes, for example, 3-aminobenzoic acid, 4-aminobenzoic acid and 5-aminopentanoic acid as well as heterocyclic
20 carboxylic acids such as tetrahydroiso-quinoline-7-carboxylic acid, 2-aminomethyl pyridine-6-carboxylic acid or the like as mentioned earlier, as replacements for the β component of the Formula (I) compounds. Thus, in a broad sense, the
25 peptidomimetics of the invention include variants for Formula (I) where β stands for the radical of a non-peptide aminoalkyl or amino-substituted aliphatic or aromatic carboxylic acid or a heterocyclic monocarboxylic acid, for example, 3-
30 aminobenzoic acid (3-ABA), 4-aminobenzoic acid (4-ABA) or 5-aminopentanoic acid (5-APA).

Other suitable β substituents which may be mentioned include those obtained by using aminomethyl- or aminocarboxylic acid derivatives
35 of other cyclic hydrophobic compounds such as furan, quinoline, pyrrole, oxazole, imidazole,

pyridine and thiazole. Generally speaking,
therefore, the β substituent may be derived from
any hydrophobic, non-peptidic aminoalkyl- or
amino-substituted aliphatic, aromatic or
5 heterocyclic monocarboxylic acid.

According to still another feature of this
embodiment of the invention, other effective
inhibitors for farnesyltransferase may be provided
by incorporating a negatively charged residue onto
10 the compounds of Formula (I). This feature of the
invention is based on a consideration of the
transition state of the farnesylation reaction and
the recognition that the functional enzyme complex
must involve a farnesyl pyrophosphate binding site
15 close to the peptide binding region. Compounds
representative of this embodiment include peptides
prepared with a phosphonate residue linked at
different distances to the cysteine sulfur. These
derivatives have been prepared by reaction of N-
20 Cbz-cysteine with ethyl 2-chloroethylphosphonate
followed by condensation with the C-terminal
methionine adduct of 4-aminobenzoic acid (or N-
deprotected VIM methyl ester). Deprotection of
the phosphonate, carboxylate and amino protecting
25 groups gives analogs (5) and (6), respectively,
which contain elements of the tetrapeptide and
farnesyl pyrophosphate residues and hence are able
to interact with binding groups in both
recognition sites in p21ras farnesyltransferase:





The above described phosphonates as contemplated herein can be structurally represented as follows:



where C, X, β and Δ are as previously described and Δ_1 is a phosphonate group joined to cysteine through the cysteine sulphur atom.

10 As indicated earlier, an important further feature of the invention is the modification of the compounds of the invention, as well as the tetrapeptide p21ras farnesyl transferase inhibitors of the formula CA_1A_2X , to provide pro-

15 drugs. This involves forming lipophilic enzyme-sensitive derivatives from the compounds by appropriately functionalizing the terminal groups. For example, the terminal amino groups and the cysteine sulfur can be reacted with benzyl


20 chloroformate to provide carbobenzyloxy ester end groups while the terminal carboxy group at the other end of the compound is converted to an alkyl or aryl ester, e.g. the methyl ester. Other

examples include alkyl esters from 1 to 10 carbons in length, activated esters such as cyanomethyl or trifluoromethyl, cholesterol, cholate or carbohydrate derivatives. The term "lipophilic", when used in this context, is meant to include, *inter alia*, methoxycarbonyl and other long chain or carbamate groups. Examples of such groups are well known to the ordinarily skilled practitioner.

Derivatization of the prior peptides CA_1A_2X and the peptidomimetics described herein with lipophilic or hydrophobic, enzyme-sensitive moieties increases the plasma membrane permeability and cellular uptake of the compounds and consequently their efficiency in inhibiting tumor cell growth.

While the carbobenzyloxy derivatives have been referred to as one way of functionalizing the peptides and peptidomimetics to improve efficiency, it will be appreciated that a variety of other groups may also be used for the purposes noted. Typical alternatives include cholesteryl, aryl or aralkyl such as benzyl, phenylethyl, phenylpropyl or naphthyl, or alkyl, typically methyl or other alkyl of, for example, up to 8 carbon atoms or more. It is contemplated that such functional groups would be attached to the cysteine sulfur and the terminal amino and carboxy groups.

Using C-ABA-M as representative of the present compounds, the functionalized pro-drug embodiment of the invention may be structurally illustrated as follows:



In the above described BBM-compounds, the "BBM" used in the formulas represents a shorthand reference to the bis-(carboxybenzyloxy)methyl esters of C β M and CVIM.

10 The functionalized derivatives of the phosphonates described earlier herein are also useful cell growth inhibitors. Correspondingly, the "BMMM" designation used with compounds refers to the carboxy benzyloxy substitution and the
15 three methyl groups in the methylated phosphoric and carboxylic acid end groups.

As noted, the purpose of the functional groups added to the parent compounds is to improve
20 entry of the compounds into tumor cells. Once in the cells, the functional groups are removed to liberate the active compound to function in its inhibitory capacity.

25 As will be recognized by those in the art, the functionalized pro-drugs of the invention can be prepared using conventional and well-known procedures for esterifying amino, SH and

carboxylic acid groups. Hence, details of such procedures are not essential for the preparation of the present pro-drugs.

EXAMPLE 1

5 SYNTHESIS OF FTI-232

A. N-BOC-4-aminobenzoic acid

4-amino-benzoic acid (10g, 72.9 mmol) was placed into a mixture of dioxane (145.8 ml) and 0.5M NaOH (145.8 ml). The solution was cooled to
10 0°C and di-t-butyl dicarbonate (23.87 g, 109.5 mmol) was added. The reaction mixture was allowed to warm to room temperature and stirred overnight. The next day, the dioxane was removed, the residue was made acidic and extracted into ethyl acetate.
15 The ethyl acetate fractions were combined and washed with 1N HCl to remove any unreacted starting material. The solution was dried over Na₂SO₄ and the solvent was removed in vacuo. The crude material was recrystallized from ethyl
20 acetate/hexanes to yield 12.2 g (70.6%) of pure product. mp 189-190°C; ¹H NMR (CD₃OD) 1.52 (9H, s), 7.49 (2H, d, J=8.6 Hz), 7.91 (2H, d, J=8.6 Hz), 9.28 (1H, s); ¹³C NMR (CD₃OD) 28.59, 81.29, 118.54, 125.30, 131.81, 145.70, 155.00, 169.80;
25 anal. calc. for C₁₂H₁₅NO₄, C: 60.76, H: 6.37, N: 5.90; found, C: 60.52, H: 6.43, N: 5.83; HRMS calc. for C₁₂H₁₅NO₄, 237.0961, found, 237.1001.

B. N-BOC-4-aminobenzoyl methionine methyl ester

Into a dried, nitrogen filled flask was
30 placed N-BOC-4-aminobenzoic acid (8.77 g, 36.97 mmol) in dry CH₂Cl₂ (148 ml) along with methionine methyl ester hydrochloride (8.12 g, 40.66 mmol). This solution was cooled in an ice bath and triethylamine (6.7 ml), EDCI (7.80 g, 40.66 mmol)
35 and hydroxybenzotriazole (HOBT, 5.50 g, 40.66

mmol) were added. The mixture was stirred overnight, diluted with more CH_2Cl_2 , and was extracted 3 times each with 1M HCl, 1M NaHCO_3 , and water. The CH_2Cl_2 was dried over MgSO_4 , and the solvent was removed in vacuo. The solid was recrystallized from ethyl acetate/ hexanes to yield 9.72 g (71.3%) of pure product. mp 184-185°C; ^1H NMR (CDCl_3) 1.53 (9H, s), 2.06-2.18 (4H, m), 2.23-2.33 (1H, m), 2.59 (2H, t, $J=7.6$ Hz), 3.80 (3H, s), 4.92 (1H, m), 7.45 (2H, d, $J=8.7$ Hz), 7.77 (2H, d, $J=8.7$ Hz); ^{13}C NMR (CDCl_3) 15.59, 28.34, 30.15, 31.64, 52.10, 52.73, 81.20, 117.73, 127.8, 128.33, 141.88, 152.33, 166.50, 172.75; anal. calc. for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_5\text{S}$, C: 56.53, H: 6.85, N: 7.29; found, C: 56.47, H: 6.86, N: 7.29; m/z (EI) 382 (M).

C. HCl-4-aminobenzoyl methionine methyl ester

N-BOC-4-aminobenzoyl methionine methyl ester (3.53 g, 9.59 mmol) was placed into CH_2Cl_2 (30-35 ml) and to it was added 3M HCl/ Et_2O (38.4 ml). After standing a white precipitate formed. After 2 hours the solution was decanted, and the crystals were collected by centrifugation. The crystals were then washed several times with fresh ether and dried overnight on the vacuum pump. Meanwhile, the filtrate was left to stand overnight to allow additional product to precipitate. The second fraction was washed with ether and dried overnight on the vacuum pump. The total yield of pure fully deprotected material was 2.87 g (93.9%) yield. mp 158-164°C; ^1H NMR (CDCl_3) 2.10 (3H, s), 2.12-2.29 (1H, m), 2.52-2.71 (1H, m), 2.59 (2H, t, $J=7.6$ Hz), 3.75 (3H, s), 4.79 (1H, m), 7.02 (2H, d, $J=8.6$ Hz), 7.55 (2H, d, $J=8.6$ Hz); ^{13}C NMR (CDCl_3) 15.23, 31.43, 31.53, 52.91, 52.43, 124.35, 130.56, 135.31, 135.76,

168.95, 173.87; HRMS calc. for $C_{13}H_{18}N_2O_3S$, 282.1038, found 282.1009.

D. N-BOC-S-trityl-cysteine-4-aminobenzoyl methionine methyl ester

5 N-BOC-S-trityl-Cys (2.86 g, 6.54 mmol) and triethylamine (1.2 ml) were placed into a dried, N_2 filled flask containing dry THF (104 ml). This was cooled to -10°C using an ice/ salt bath and isobutyl chloroformate (0.9 ml), IBCF, was added.

10 The solution was stirred at -10°C for 40 minutes and HCl-4-aminobenzoyl methionine methyl ester (2.08 g, 6.54 mmol) in dry CH_2Cl_2 (34.1 ml) with triethylamine (1.2 ml, 1.3 eq) was added. The solution warmed to room temperature and was

15 stirred overnight under N_2 . The solvent was then removed in vacuo and the residue was taken up in CH_2Cl_2 and extracted several times each with 1M HCl, H_2O and brine (saturated NaCl). The organic layer was dried over Na_2SO_4 and the solvent was

20 removed in vacuo. The pale yellow foam was then chromatographed on silica gel using a 2:1 hexanes, ethyl acetate elution mixture to yield 2.62 g (54.9%) of pure product. mp $110-111^\circ\text{C}$; $[\alpha]^{25}_D = -8.0^\circ$ ($c=1$, CH_3OH); ^1H NMR (CDCl_3) 1.44 (9H, s),

25 2.11-2.18 (4H, m), 2.22-2.34 (1H, m), 2.59 (2H, t, $J=7.4$ Hz), 2.66-2.83 (2H, m), 3.80 (3H, s), 3.98 (1H, m), 4.84 (1H, m), 4.92 (1H, m), 6.96 (1H, d, $J=7.7$ Hz), 7.23-7.33 (9H, m), 7.43-7.46 (6H, m), 7.51 (2H, d, $J=8.5$ Hz), 7.74 (2H, d, $J=8.5$ Hz),

30 8.51 (1H, s); ^{13}C NMR (CDCl_3) 15.53, 28.34, 30.72, 30.89, 33.60, 52.23, 52.88, 54.95, 60.50, 67.13, 80.64, 118.81, 119.31, 126.94, 128.07, 128.30, 129.53, 141.06, 144.38, 156.31, 167.02, 170.13, 174.49; anal calc for $\text{C}_{40}\text{H}_{44}\text{N}_3\text{O}_6\text{S}_2 \cdot \text{H}_2\text{O}$, C: 64.50, H: 6.22, N: 5.64; found C: 64.14 H: 6.19, N: 5.56.

35

E. HCl-cysteine-4-aminobenzoyl methionine methyl ester

N-BOC-S-trityl-cysteine-4-aminobenzoyl methionine methyl ester (1 g, 1.37 mmol) was placed into a flask and taken up in CH₃OH (13.7 ml). To this solution was added a solution of mercuric chloride (0.75 g, 2.74 mmol) in CH₃OH (13.7 ml). Upon addition of the mercuric chloride, a white precipitate began to form. The mixture was heated on a steam bath at 65°C for 35 minutes and then it was cooled and the precipitate was filtered and washed sparingly with cold CH₃OH. After drying for several minutes on the filter, the solid was placed into a 50 ml 3-neck flask fitted with a gas inlet and outlet. Approximately 20-30 ml of CH₃OH was added and H₂S gas was bubbled through the heterogeneous solution for 30 minutes. Upon addition of the gas, the white solution turned orange and then black. The solution was centrifuged and the clear, colorless liquid was dried to give a white foam. This solid was placed on the vacuum pump for a short period and then was taken up in CH₂Cl₂ (10 ml) and the product was precipitated with a 3-4M HCl/ Et₂O solution. The precipitate was collected by centrifugation and was washed with ether until pH was neutral. After drying under vacuum overnight, 0.38 g (66.5%) of product was obtained that was >95% pure by HPLC. mp foamed 141-143°C, decomp 195°C; $[\alpha]^{25}_D = +3^\circ$ (c=1, H₂O); ¹H NMR (CD₃OD) 2.09 (3H, s), 2.14-2.26 (1H, m), 2.51-2.67 (3H, m), 3.05 (1H, dd, J=14.8 Hz, 7.3 Hz), 3.17 (1H, dd, J=14.8 Hz, 4.8 Hz), 3.74 (3H, s), 4.17 (1H, J=7.3 Hz, 4.8 Hz), 4.75-4.81 (1H, m), 7.74 (2H, d, J=8.6 Hz), 7.87 (2H, d, J=8.6 Hz), 8.67 (1H, d, J=8.4 Hz); ¹³C NMR (CD₃OD) 15.23, 26.38, 31.43, 31.56, 52.88, 53.30, 56.92, 120.46, 129.58, 130.75, 142.33, 166.91, 169.66,

174.06; anal calc for $C_{16}H_{24}ClN_3O_4S_2$, C: 45.55 H: 5.73, N: 9.96; found C: 45.31, H: 5.84, N: 9.79.

F. HCl-cysteine-4-aminobenzoyl methionine FTI-232

HCl-cysteine-4-aminobenzoyl methionine methyl ester (0.51 g, 0.7 mmol) was taken up in THF (4.1 ml) and to this solution was added 0.5 M LiOH (2.9 ml) at 0°C. The heterogeneous solution was stirred at 0°C for 35-40 minutes and then the THF was removed in vacuo. The residue was taken up in CH_2Cl_2 , and was washed three times with 1M HCl followed by brine. The organic solution was dried over Na_2SO_4 and the solvent was removed in vacuo. The pale yellow solid was taken up in 3 ml of CH_2Cl_2 and the product was precipitated with 3-4 M HCl/ Et_2O . The solid was collected by centrifugation, washed several times with ether until the ether washings were neutral and the process repeated until the HPLC appeared pure. A final yield of 78.6 mg (27.5%) of pure product was obtained. mp sub 157°C, decomp 211°C; $[\alpha]^{25}_D = +10^\circ$ (c=0.8, H_2O); 1H NMR (CD_3OD) 2.09 (3H, s), 2.17-2.32 (1H, m), 2.53-2.66 (3H, m), 3.06 (1H, dd, J=14.6 Hz, 7.2 Hz), 3.19 (1H, dd, J=14.6 Hz, 4.6 Hz), 4.21 (1H, dd, J=7.23 Hz, 4.63 Hz), 4.73-4.78 (1H, m), 7.75 (2H, d, J=8.1 Hz), 7.87 (2H, d, J=8.1 Hz); ^{13}C NMR (CD_3OD) 15.23, 26.33, 31.58, 31.86, 53.24, 56.98, 120.48, 129.59, 131.10, 142.26, 166.89, 169.66, 175.29; anal calc for $C_{15}H_{22}ClN_3O_4S_2$, C: 44.16, H: 5.44, N: 10.30; found C: 45.45, H: 5.62, N: 10.03; m/z (FAB) for free amine, 371 ($M+1$).

EXAMPLE 2

SYNTHESIS OF FTI-260**A. N-BOC-4-amino-3-methylbenzoic acid**

4-amino-3-methylbenzoic acid (5 g, 33.1 mmol)
5 was reacted according to the same procedure as N-BOC-4-aminobenzoic acid. The orange-brown solid was recrystallized from ethyl acetate and hexanes to yield 4.99 g (60%) of tan prismatic crystals.
mp 180-182°C; ¹H NMR (CD₃OD) 1.51 (9H, s), 2.27
10 (3H, s), 7.66 (1H, d, J=8.1 Hz), 7.79-7.82 (2H, m), 8.32 (1H, s); ¹³C NMR (CD₃OD) 17.98, 28.62, 81.47, 123.12, 127.05, 129.14, 130.65, 132.99, 142.45, 155.33, 168.70; anal calc for C₁₃H₁₁NO₄, C: 62.15, H: 6.82, N: 5.58; found C: 62.07, H: 6.86,
15 N: 5.46; m/z (EI) 251; HRMS calc. for C₁₃H₁₁NO₄, 251.1158; found, 251.1153.

B. N-BOC-4-amino-3-methylbenzoyl methionine methyl ester

N-BOC-4-amino-3-methylbenzoic acid (2.00 g,
20 7.96 mmol) was reacted with methionine methyl ester hydrochloride (1.75 g, 8.76 mmol), EDCI (1.68 g, 8.76 mmol), HOBT (1.18 g, 8.76 mmol) and Et₃N (1.4 ml) in dry CH₂Cl₂ (31.8 ml) according to the procedure described for N-BOC-4-aminobenzoyl
25 methionine methyl ester in Example 1. The crude material was recrystallized from ethyl acetate and hexanes to yield 2.61 g (85.7%) of pure product.
mp 163-165°C; ¹H NMR (CDCl₃) 1.54 (9H, s), 2.06-2.18
(4H, m), 2.23-2.34 (4H, m), 2.59 (2H, t, J=6.8
30 Hz), 3.80 (3H, s), 4.92 (1H, m), 6.45 (1H, s), 6.88 (1H, d, J=7.5 Hz), 7.63 (1H, d, J=8.6 Hz), 7.66 (1H, s), 8.05 (1H, d, J=8.6); ¹³C NMR (CDCl₃)
15.47, 17.61, 28.22, 30.03, 31.55, 51.93, 52.57, 81.04, 118.73, 125.62, 127.66, 129.54, 139.89,
35 152.34, 166.58, 172.66.

C. HCl-4-amino-3-methylbenzoyl methionine methyl ester

N-BOC-4-amino-3-methylbenzoyl methionine methyl ester (0.99 g, 2.59 mmol) was dissolved in CH₂Cl₂ (15-20 ml) and precipitated with 3M HCl/Et₂O (20.7 ml). 0.83 g (96.6%) of pale orange precipitate was obtained after drying overnight on the vacuum pump. mp 157-159°C; ¹H NMR (CD₃OD) 2.04 (3H, s), 2.11-2.25 (1H, m), 2.47 (3H, s), 2.52-2.68 (3H, m), 3.74 (3H, s), 4.75-4.80 (1H, m), 7.48 (1H, d, J=8.2 Hz), 7.81 (2H, d, J=8.2 Hz), 7.87 (1H, s); ¹³C NMR (CD₃OD) 15.23, 17.28, 31.43, 31.51, 52.91, 53.37, 124.41, 127.85, 131.99, 133.63, 134.14, 135.65, 169.05, 173.84; anal. calc. for C₁₄H₂₁N₂O₃S, C: 50.52, H: 6.36, N: 8.42; found C: 50.71, H: 6.40, N: 8.34.

D. N-BOC-S-trityl-cysteine-4-amino-3-methylbenzoyl methionine methyl ester

N-BOC-S-trityl-cysteine (0.55 g, 1.25 mmol) in dry THF (25 ml) was reacted with Et₃N (0.19 ml), IBCF (0.16 ml, 1.25 mmol) at -10 °C as described above. HCl-4-amino-3-methylbenzoyl methionine methyl ester (0.42 g, 1.25 mmol) in dry CH₂Cl₂ (6.5 ml) with Et₃N (0.26 ml) was added at -10°C and the reaction mixture was allowed to stir overnight under nitrogen. Workup was carried out as described above and the crude material was chromatographed on silica gel using a 2:1 mixture of hexanes and ethyl acetate as an elution mixture to give 0.12 g (13.9%) of pure product. mp 83-85°C; [α]_D²⁵ = -14.0° (c=1, CH₃OH); ¹H NMR (CDCl₃) 1.44 (9H, s), 2.10-2.17 (4H, m), 2.22-2.32 (4H, m), 2.61 (2H, t, J=6.57 Hz), 2.68-2.70 (1H, m), 2.85-2.90 (1H, m), 3.79 (3H, s), 3.93-4.08 (1H, s), 4.84-4.88 (1H, m), 4.90-4.95 (1H, m), 6.95 (1H, d, J=7.00 Hz), 7.20-7.33 (9H, m), 7.39 (1H, d, J=6.96 Hz),

7.44-7.47 (6H,m), 7.59 (1H, d, J=8.46 Hz), 7.65 (1H, s), 8.12 (1H,d, J=8.22 Hz), 8.31 (1H,s); ¹³C NMR (CDCl₃) 15.39 17.55, 27.70, 28.17, 30.00, 31.43, 31.41, 51.90, 52.51, 59.95, 67.30, 80.74, 84.54, 120.74, 125.33, 126.70, 126.83, 127.89, 128.00, 129.40, 138.92, 144.22, 166.50, 166.89, 168.87, 172.56.

E. TFA-cysteine-4-amino-3-methylbenzoyl methionine FTI-260

10 N-BOC-S-trityl-cysteine-4-amino-3-methylbenzoyl methionine methyl ester (0.27 g, 0.37 mmol) in THF (2.1 ml) was deprotected with 0.5M LiOH (2.9 ml) over 1.5 h at room temperature. The solvent was removed in vacuo and the residue
15 was taken up in CH₂Cl₂ and extracted 3 times with 1N HCl followed by extraction with brine. The organic solution was dried over Na₂SO₄ and the solvent was removed in vacuo to give 0.19 g (73.5 %) of the free acid. The free acid was then taken
20 up in CH₂Cl₂ (1.4 ml) and Et₃SiH (0.04 ml) was added followed by trifluoroacetic acid, TFA (1.4 ml). The reaction mixture was stirred at room temperature for 1 hour. The TFA was removed and the residue was dissolved in H₂O and extracted with
25 Et₂O until all of the trityl derivative was removed. The water was lyophilized and a crude HPLC showed that the material was impure and contained diastereomers. The product was purified on the preparative HPLC using 0.1% TFA in water
30 and acetonitrile elution mixture to give 2 diastereomers and only the major component (determined according to the major compound in the HPLC trace) was characterized. mp sub 112°C, foamed 158-163°C, decomp 196-197°C; [α]²⁵_D=+12.7° (c=0.6 H₂O), [α]²⁵_D=+ 21.0° (c=1 H₂O); ¹H NMR (CD₃OD)
35 2.09-2.17 (4H, m), 2.19-2.30 (1H, m), 2.36 (3H,

s), 2.57-2.65 (2H, m), 3.08 (1H, dd, J=14.6 Hz, 6.9 Hz), 3.19 (1H, dd, J=14.6 Hz, 5.2 Hz), 4.25 (1H, dd, J=6.9, 5.2 Hz), 4.70-4.75 (1H, m), 7.64 (1H, d, J=8.4 Hz), 7.69-7.73 (1H, m), 7.77 (1H, s); ¹³C NMR (CD₃OD) 15.23, 18.28, 26.54, 31.58, 32.06, 53.53, 56.66, 125.54, 125.77, 126.74, 131.04, 133.24, 139.26, 167.53, 169.70, 175.59.

EXAMPLE 3

SYNTHESIS OF FTI-261

10 A. N-BOC-4-amino-3-methoxybenzoic acid

4-amino-3-methoxybenzoic acid (1 g, 5.98 mmol) was reacted with di-t-butyl dicarbonate (1.96 g, 6.58 mmol) in dioxane (12 ml) and 0.5 M NaOH (12 ml) according to the same procedure as N-BOC-4-aminobenzoic acid. 1.50 g (93.7%) of tan crystals were obtained after recrystallization from ethyl acetate and hexanes. mp 176-178°C; ¹H NMR (CD₃OD) 1.52 (9H, s), 3.92 (3H, s), 7.56 (1H, s), 7.62 (1H, d, J=8.4 Hz), 7.96 (1H, s), 8.03 (1H, d, J=8.4 Hz); ¹³C NMR (CD₃OD) 28.53, 56.35, 81.78, 112.01, 118.58, 124.20, 125.76, 133.84, 149.04, 154.20, 169.60; HRMS calc. for C₁₃H₁₁NO₅, 267.1107; found, 267.1103.

25 B. N-BOC-4-amino-3-methoxybenzoyl methionine methyl ester

N-BOC-4-amino-3-methoxybenzoic acid (0.35 g, 1.31 mmol) was reacted with methionine methyl ester hydrochloride (0.9 g, 1.43 mmol) using EDCI as in N-BOC-4-aminobenzoyl methionine methyl ester. After recrystallization from ethyl acetate and hexanes, 0.36 g (57.2 %) of pure product was obtained. mp 163-165°C; ¹H NMR (CDCl₃) 1.53 (9H, s), 2.09-2.18 (4H, m), 2.23-2.35 (1H, m), 2.60 (2H, t, J=6.9 Hz), 3.80 (3H, s), 3.93 (3H, s), 4.92 (1H, br s), 6.93 (1H, d, J=7.6 Hz), 7.25 (1H,

m), 7.31 (1H, d, J=10.2 Hz), 7.44 (1H, s), 8.15 (1H, d, J=8.5 Hz); ¹³C NMR (CDCl₃) 15.47, 28.23, 30.09, 31.48, 52.06, 52.54, 55.81, 80.82, 98.06, 109.38, 116.66, 119.31, 131.52, 147.23, 152.31, 166.57, 172.58; m/z (FAB) 413 (M+1).

C. HCl-4-amino-3-methoxybenzoyl methionine methyl ester

N-BOC-4-amino-3-methoxybenzoyl methionine methyl ester (0.71 g, 1.79 mmol) was taken up in CH₂Cl₂ (4 ml) and precipitated with 3-4M HCl/ Et₂O (12 ml). The precipitate was washed as usual with Et₂O and dried overnight under vacuum to result in 0.55 g (88.3%) of reddish material. mp 176-177°C; ¹H NMR (CD₃OD) 2.08 (3H, s), 2.21 (2H, m), 2.61 (2H, m), 3.74 (3H, s), 4.02 (3H, s), 4.79 (1H, m), 7.50 (1H, d, J=8.2 Hz), 7.57 (1H, d, J=4.1 Hz), 7.67 (1H, s); ¹³C NMR (CD₃OD) 15.26, 31.34, 31.42, 52.95, 53.38, 57.12, 112.29, 121.43, 124.57, 124.77, 136.15, 153.67, 168.79, 173.81.

D. N-BOC-S-trityl-cysteine-4-amino-3-methoxybenzoyl methionine methyl ester

N-BOC-S-trityl-cysteine (0.76 g, 1.74 mmol) in dry THF (27.5 ml) was reacted with Et₃N (0.24 ml), IBCF (0.23 ml, 1.74 mmol) at -10°C as described above. HCl-4-amino-3-methoxybenzoyl methionine methyl ester (0.55 g, 1.58 mmol) in dry CH₂Cl₂ (8.7 ml) with Et₃N (0.30 ml) was added to the mixture and was allowed to stir overnight under nitrogen. It was worked up as described for N-BOC-S-trityl-cysteine-4-aminobenzoyl methionine methyl ester in Example 1, and the crude material was chromatographed on silica gel using a 2:1 mixture of hexanes and ethyl acetate to give 0.18 g (15.2%) of pure product. ¹H NMR (CDCl₃) 1.45 (9H, s), 2.05-2.33 (5H, m), 2.57-2.65 (2H, m),

2.68-2.72 (1H, m), 2.75-2.96 (1H, m), 3.78 (3H, s), 3.84 (3H, s), 4.90-5.00 (1H, m), 5.03-5.18 (1H, m), 7.17-7.48 (17H, m), 8.30-8.38 (1H, m), 8.65 (1H, br s).

5 **E. TFA-Cysteine-4-amino-3-methoxybenzoyl methionine FTI-261**

 N-BOC-S-trityl-cysteine-4-amino-3-methoxybenzoyl methionine methyl ester (0.18 g, 0.24 mmol) was deprotected with LiOH at room
10 temperature as described above to give the free acid. The free acid was then further deprotected in CH₂Cl₂ (1.2 ml) with Et₃SiH (0.04 ml, 0.24 mmol) and TFA (1.2 ml). The product was worked up as
15 described for HCl-cysteine-4-aminobenzoyl methionine in Example 1, and HPLC revealed that the product was impure. The crude material was then purified on the HPLC using 0.1% TFA in water and acetonitrile as eluting solvents to result in
20 two pure samples that were expected to be diastereomers. The major component (determined according to the major compound in the HPLC trace) was characterized as follows. mp sub 109°C, decomp 191-193°C; $[\alpha]^{25}_D = -30.0^\circ$ (c=1, H₂O), $[\alpha]^{25}_D = +19.0^\circ$ (c=1, H₂O); ¹H NMR (CD₃OD) 2.10 (3H, s), 2.12-2.18 (1H, m), 2.20-2.32 (1H, m), 2.53-2.71 (2H, m), 3.00 (1H, dd, J=14.6, 7.5), 3.15 (1H, dd, J=14.58, 4.8), 4.77 (1H, dd, J=7.5, 4.8), 7.50 (1H, d, J=8.4 Hz), 7.56 (1H, s), 8.23 (1H, d, J=8.4 Hz); ¹³C NMR (CD₃OD) 15.20, 26.65, 31.60,
25 31.76, 53.27, 56.58, 56.76, 111.04, 121.08, 122.14, 130.85, 131.85, 150.88, 167.21, 169.61, 175.36; m/z (FAB) for free amine, 402 (M+1).
30

EXAMPLE 4

SYNTHESIS OF FTI-272**A. 4-nitro-2-phenyltoluene**

2-bromo-4-nitrotoluene (2.16 g, 10.00 mmol) and phenyl boric acid (1.46 g, 12.00 mmol) were dissolved into anhydrous DMF (25 ml) under nitrogen. To this mixture was added $\text{Pd}(\text{Ph}_3\text{P})_4$ (0.58 g, 5%). The mixture was heated at 100 °C overnight. The solution was poured onto 1N HCl and extracted with Et_2O . The crude material was chromatographed on silica gel using hexanes as an eluent. After recrystallization from ethanol, 1.23 g (57.6%) of pale orange needles were obtained. mp 69 - 71°C; ^1H NMR (CDCl_3) 2.36 (3H, s), 7.29-7.40 (2H, m), 7.41-7.49 (5H, m), 8.07-8.10 (2H, m); ^{13}C NMR (CDCl_3) 20.68, 121.96, 124.51, 127.78, 128.41, 128.83, 131.06, 139.44, 142.97, 143.48, 146.05; anal calc. for $\text{C}_{13}\text{H}_{11}\text{NO}_2$, C:73.26, H:5.20, N:6.57; found, C:73.10, H:5.12, N:6.50; m/z (EI) 213; HRMS calc. for $\text{C}_{13}\text{H}_{11}\text{NO}_2$, 213.0790; found, 213.0793.

B. 4-nitro-2-phenylbenzoic acid

4-nitro-2-phenyltoluene (0.50 g, 2.34 mmol) was dissolved in water (4.6 ml) and pyridine (2.3 ml). The mixture was heated to reflux and KMnO_4 (1.85 g, 11.70 mmol) was added. The reaction mixture was heated overnight and the solution was filtered and washed several times with boiling water. The aqueous solution was made acidic and the product was extracted into ethyl acetate. The ethyl acetate was dried over Na_2SO_4 and the solvent removed in vacuo to result in 0.37 g (67.9%) of pure yellow product. mp 174-176°C; ^1H NMR (CD_3OD) 7.38-7.48 (5H, m), 7.96 (1H, d, $J=8.5$ Hz), 8.21 (1H, d, $J=2.3$ Hz), 8.28 (1H, dd, $J=8.48$, 2.37); ^{13}C NMR (CD_3OD) 122.95, 126.09, 129.27, 129.42, 129.49,

131.56, 139.26, 140.42, 144.41, 150.17, 170.52; m/z (EI) 243 (M).

C. 4-nitro-2-phenylbenzoyl methionine methyl ester

5 4-nitro-2-phenylbenzoic acid (0.30 g, 1.23 mmol), methionine methyl ester hydrochloride salt (0.27 g, 1.35 mmol), EDCI (0.26 g, 1.35 mmol), HOBT (0.18 g, 1.35 mmol) and Et₃N (0.19 ml) in dry CH₂Cl₂ (4.9 ml) were reacted according to the above
10 procedure and worked up as described for N-BOC-4-aminobenzoyl methionine methyl ester in Example 1. After recrystallization from ethyl acetate and hexanes, 0.41 g (85.5%) of pure product was isolated. mp 98-101°C; ¹H NMR (CDCl₃) 1.62-1.73
15 (1H, m), 1.79-1.88 (1H, m), 1.91 (3H, s), 1.99 (2H, t, J=7.2 Hz), 3.59 (3H, s), 4.53 (1H, m), 6.45 (1H, d, J=7.8 Hz), 7.33-7.40 (5H, m), 7.67 (1H, d, J=8.3 Hz), 8.07-8.12 (2H, m); ¹³C NMR (CDCl₃) 14.92, 29.11, 30.67, 51.51, 52.29, 121.86,
20 124.74, 128.27, 128.60, 128.69, 129.52, 137.50, 140.56, 141.02, 148.09, 167.23, 171.23; m/z (FAB), 389 (M+1).

D. 4-amino-2-phenylbenzoyl methionine methyl ester

25 4-nitro-2-phenylbenzoyl methionine methyl ester (0.35 g, 0.90 mmol) was taken up in ethyl acetate (9.0 ml). To this mixture was added SnCl₂·2H₂O (1.02 g, 4.50 mmol) and the reaction was heated under nitrogen at reflux for 1h. The
30 mixture was poured onto ice, the solution was made basic using NaHCO₃ and the product was extracted into ethyl acetate several times (7-8). The ethyl acetate fractions were combined washed with brine and dried over Na₂SO₄ and the solvent was removed
35 in vacuo to give 0.24 g (73.4%) of yellow solid.

¹H NMR (CDCl₃) 1.58-1.70 (1H, m), 1.80-1.92 (1H, m), 1.98 (3H, s), 2.06 (2H, t, J=7.7 Hz), 3.62 (3H, s), 4.00 (2H, br s), 4.56-4.63 (1H, m), 5.84 (1H, d, J=7.7 Hz), 6.50 (1H, s), 6.61 (1H, d, J=8.4 Hz), 7.29-7.42 (5H, m), 7.58 (1H, d, J=8.3 Hz); ¹³C NMR (CDCl₃) 15.02, 29.25, 31.25, 51.57, 52.15, 113.27, 115.88, 123.52, 127.56, 128.37, 128.44, 130.92, 140.66, 141.44, 148.53, 168.58, 171.91.

10 **E. N-BOC-S-trityl-cysteine-4-amino-2-phenylbenzoyl methionine methyl ester**

N-BOC-S-trityl-cysteine (0.31 g, 0.66 mmol) in dry THF (11 ml) was reacted with Et₃N (0.10 ml), IBCF (0.09 ml, 0.73 mmol) at -10 °C as described for N-BOC-S-trityl-cysteine-4-aminobenzoyl methionine methyl ester in Example 1. 4-amino-2-phenylbenzoyl methionine methyl ester (0.24g, 0.66 mmol) in dry CH₂Cl₂ (3.5 ml) was added and the mixture was allowed to stir overnight under nitrogen. It was worked up as described as further described for N-BOC-S-trityl-cysteine-4-aminobenzoyl methionine methyl ester in Example 1, and after drying the crude material was chromatographed on silica gel using a 2:1 mixture of hexanes and ethyl acetate to give 84.70 mg (16.0%) of pure product. mp 100-103°C; ¹H NMR (CDCl₃) 1.41 (9H, s), 1.61-1.78 (1H, m), 1.84-1.95 (1H, m), 2.00 (3H, s), 2.05 (2H, t, J=7.6 Hz), 2.63 (1H, dd, J=12.7 Hz, 6.9 Hz), 2.72 (1H, dd, J=12.7 Hz, 5.51 Hz), 3.64 (3H, s), 4.02 (1H, br s), 4.58-4.63 (1H, m), 4.90 (1H, d, J=7.4 Hz), 6.10 (1H, d, J=6.6 Hz), 7.18-7.30 (10H, m), 7.37-7.44 (11H, m), 7.50 (1H, s), 7.58 (1H, d, J=8.2 Hz), 8.69 (1H, s); ¹³C NMR (CDCl₃) 15.21, 28.20, 29.38, 31.24, 33.00, 51.77, 52.35, 54.15, 67.30, 80.85, 118.18, 120.86, 126.88, 127.90, 128.03,

128.56, 128.66, 129.44, 129.79, 130.14, 156.00,
168.52, 169.11, 171.85.

**F. TFA·Cysteine-4-amino-2-phenylbenzoyl
methionine FTI-272**

5 N-BOC-S-trityl-cysteine-4-amino-2-
phenylbenzoyl methionine methyl ester (84.70 mg,
0.11 mmol) of was taken up in THF (0.62 ml) and to
this was added 0.5 M LiOH (0.32 ml) at 0 °C. The
mixture was stirred at 0 °C for 35 minutes. The
10 solvent was removed in vacuo using a cold water
bath on the rotovap. The residue was worked up as
described for HCl-cysteine-4-aminobenzoyl
methionine in Example 1, and 60 mg of the free
acid was obtained. This was then dissolved into
15 CH₂Cl₂ (0.8 ml) and Et₃SiH (0.01 ml) was added
followed by TFA (0.8 ml). The reaction mixture
was stirred at room temperature for 1 h and worked
up as described for TFA·cysteine-4-amino-3-
methylbenzoyl methionine in Example 2. After
20 lyophilization, 0.0387 g (84.0%) was obtained.
HPLC revealed that no epimerization had occurred,
however the material was purified on the HPLC to
eliminate baseline impurities. mp 150-154°C;
[α]²⁵_D=+21.5° (c=0.7, H₂O/ CH₃OH); ¹H NMR (CD₃OD)
25 1.62-1.79 (1H, m), 2.00-2.10 (5H, m), 2.16-2.18
(1H, m), 3.03 (1H, dd, J=14.7 Hz, 7.3 Hz), 3.15
(1H, dd, J=14.7 Hz, 4.8 Hz), 4.46 (1H, br s),
7.37-7.41 (5H, m), 7.52- 7.55 (1H, m), 7.65-7.67
(2H, m); ¹³C NMR (CD₃OD) 15.03, 26.35, 31.78,
30 32.79, 57.01, 119.40, 122.35, 128.95, 129.62,
129.71, 130.15, 133.49, 140.50, 141.36, 142.53,
167.05, 167.76, 172.51; anal. calc. for
C₂₃H₂₆F₃N₃O₆S₂, C: 49.20, H: 4.67, N: 7.48; found, C:
49.14 H: 4.71, N: 7.42.

EXAMPLE 5

**HCl·cysteine-4-amino-2-phenylbenzoyl methionine
methyl ester FTI274**

N-BOC-S-trityl-cysteine-4-amino-2-phenylbenzoyl methionine methyl ester (0.06 g, 0.075 mmol) was dissolved into methanol (2 ml) and to it was added HgCl₂ (0.04 g) in methanol (1 ml). The reaction was carried out as described above to yield 15.7 mg of slightly impure compound by HPLC. mp 130-132°C; ¹H NMR (CD₃OD) 1.72-1.84 (1H, m), 1.90-2.24 (6H, m), 3.05 (1H, dd, J=14.6 Hz, 8.5 Hz), 3.19 (1H, dd, J=14.6 Hz, 3.6 Hz), 3.69 (3H, s), 4.22 (1H, dd, J=.5 Hz, 3.6 Hz), 4.48-4.53 (1H, m), 7.33-7.43 (5H, m), 7.51 (1H, d, J=8.9 Hz), 7.70-7.72 (2H, m); ¹³C NMR (CD₃OD) 15.04, 26.36, 30.88, 31.36, 52.85, 53.05, 56.93, 119.42, 122.38, 128.88, 129.55, 129.73, 130.05, 133.17, 140.55, 141.32, 142.52, 166.92, 172.61, 173.58; anal. calc. for C₂₄H₂₉ClN₃O₆S₂·2H₂O, C: 51.20, H: 5.86, N: 8.14; found, C: 51.23 H: 5.60, N: 8.22.

EXAMPLE 6

SYNTHESIS OF FTI-275**A. 2-bromo-4-nitrobenzoic acid**

2-bromo-4-nitrotoluene (5.00 g, 23.14 mmol) was dissolved into pyridine (23 ml) and water (46 ml). The heterogeneous mixture was heated to 60 °C and KMnO₄ (18.29 g, 115.7 mmol) was added carefully. The mixture was then heated under reflux overnight. The reaction mixture was filtered and washed with boiling water. The solution was then made acidic and extracted into ethyl acetate, dried over Na₂SO₄, and the solvent was removed in vacuo. A crude NMR revealed remaining starting material so the solid was taken up in NaOH and washed with hexanes. The aqueous phase was made acidic and the product was

extracted into ethyl acetate. The ethyl acetate fractions were combined and dried over Na_2SO_4 and the solvent was removed in vacuo to yield 3.72 g (65.4%). mp 158-160°C; ^1H NMR (CD_3OD) 7.81 (1H, d, J=8.5 Hz), 8.08 (1H, d, J=8.5 Hz), 8.30 (1H, s); ^{13}C NMR (CD_3OD) 121.96, 122.75, 129.36, 132.24, 139.52, 149.54, 167.75; anal. calc. for $\text{C}_7\text{H}_4\text{BrNO}_4$ + 0.1 ethyl acetate, C: 34.88, H: 1.90, N: 5.50; found, C: 34.68, H: 1.86, N: 5.82.

10 B. 3,5-dimethylphenyl boronic acid

Mg turnings (1.44 g, 59.43 mmol) were covered with dry THF (18.8 ml) in a dried, N_2 filled flask fitted with an addition funnel and reflux condenser. To this was added 5-bromo-m-xylene (10 g, 54.03 mmol) in THF (15 ml) after initiation of the Grignard reaction. The addition was carried out over several minutes and the reaction mixture was heated at reflux for 1-2 h until most of the Mg had reacted. The reaction mixture was then cooled and transferred to an addition funnel fitted to a N_2 filled flask containing triisopropyl borate (24.9 ml) at -70 °C. The dropwise addition was carried out over several minutes and the mixture warmed to room temperature and stirred overnight. The grey solution was poured onto 2 M HCl and immediately turned yellow. The solution was extracted into Et_2O and the Et_2O fractions were combined, dried over MgSO_4 and the solvent was removed in vacuo to yield 2.41 g (29.7%). mp 249-251°C; ^1H NMR (CDCl_3) 2.44 (6H, s), 7.23 (1H, s), 7.84 (2H, s); ^{13}C NMR (CD_3OD) 21.36, 133.28, 134.39, 137.48.

C. 4-nitro-2-(3,5-dimethylphenyl)benzoic acid

2-bromo-4-nitrobenzoic acid (0.50 g, 2.03 mmol) and 3,5-dimethylphenyl boronic acid (0.34 g,

2.23 mmol) were dissolved into anhydrous DMF (dimethylformamide) (25 ml) under nitrogen. To this mixture was added Cs_2CO_3 (1.66 g, 5.08 mmol) followed by $\text{Pd}(\text{Ph}_3\text{P})_4$ (0.12 g, 5%). The mixture was heated at 100°C overnight. The solution was poured onto 1N HCl and extracted into Et_2O . It was dried over MgSO_4 and the solvent was removed in vacuo. The crude material was chromatographed on silica gel using a 9:1 mixture of hexanes and ethyl acetate to yield 0.34 g (61.7%) of pure product. ^1H NMR (CDCl_3) 2.36 (6H, s), 6.99 (2H, s), 7.07 (1H, s), 8.03 (1H, d, $J=9.0$ Hz), 8.23-8.25 (2H, m); ^{13}C NMR (CDCl_3) 21.28, 121.68, 123.68, 125.74, 126.07, 130.22, 131.19, 131.31, 135.04, 138.21, 144.74, 170.75.

D. 4-nitro-2-(3,5-dimethylphenyl)benzoyl methionine methyl ester

4-nitro-2-(3,5-dimethylphenyl)benzoic acid (0.15 g, 0.55 mmol), methionine methyl ester hydrochloride salt (0.11 g, 0.55 mmol), EDCI (0.11 g, 0.55 mmol), HOBT (0.07 g, 0.55 mmol) and Et_3N (0.08 ml) in dry CH_2Cl_2 (2.2 ml) were reacted and worked up according to the procedure for N-BOC-4-aminobenzoyl methionine methyl ester in Example 1. After recrystallization from ethyl acetate and hexanes, 0.13 g (58.4%) of pure product was isolated. mp 122-124°C; ^1H NMR (CDCl_3) 1.2-1.84 (1H, m), 1.85-1.97 (1H, m), 2.01 (3H, s), 2.05 (3H, t, $J=7.7$ Hz), 2.38 (6H, s), 3.70 (3H, s), 4.67-4.74 (1H, m), 6.03 (1H, d, $J=7.9$ Hz), 7.05 (2H, s), 7.09 (1H, s), 7.84-7.87 (1H, m), 7.84-7.87 (1H, m), 8.23-8.26 (2H, m); ^{13}C NMR (CDCl_3) 15.20, 21.26, 29.22, 31.15, 51.79, 52.57, 122.07, 25.11, 126.27, 130.03, 130.53, 137.77, 138.82, 140.29, 141.56, 148.41, 167.14, 171.53.

E. 4-amino-2-(3,5-dimethylphenyl)benzoyl methionine methyl ester

4-nitro-2-(3,5-dimethylphenyl)benzoyl methionine methyl ester (0.11 g, 0.26 mmol) was taken up in ethyl acetate (3.0 ml). To this mixture was added $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (0.30 g, 1.30 mmol) and the reaction was heated under nitrogen at reflux for 6h. The mixture was worked up as described for 4-amino-2-phenylbenzoyl methionine methyl ester in Example 2, to give 0.15 g of a yellow film that was wet with solvent. The material was otherwise pure by NMR and was used without further purification. ^1H NMR (CDCl_3) 1.60-1.70 (1H, m), 1.80-1.90 (1H, m), 1.99 (3H, s), 2.05 (2H, t, $J=7.6$ Hz), 2.33 (6H, s), 3.64 (3H, s), 3.93 (2H, br s), 4.61-4.64 (1H, m), 5.82 (1H, d, $J=7.7$ Hz), 6.49 (1H, d, $J=2.3$ Hz), 6.62 (1H, dd, $J=8.4$ Hz, 2.4 Hz), 6.98 (2H, s), 7.00 (1H, s), 7.65 (1H, d, $J=8.3$ Hz); ^{13}C NMR (CDCl_3) 15.08, 21.17, 29.28, 31.49, 51.70, 52.18, 113.30, 115.94, 123.55, 126.36, 129.32, 131.23, 138.15, 140.72, 141.92, 148.40, 168.45, 172.01.

F. N-BOC-S-trityl-cysteine-4-amino-2-(3,5-dimethylphenyl)benzoyl methionine methyl ester

4-amino-2-(3,5-dimethylphenyl)benzoyl methionine methyl ester (0.10g, 0.26 mmol) was dissolved into dry CH_2Cl_2 (1.4 ml) and it was allowed to stand. In another flask, N-BOC-S-trityl-Cys (0.12 g, 0.26 mmol) was dissolved into THF (4.4 ml) and was reacted with IBCF (0.03 ml) and Et_3N (0.04 ml) as described above. The product was worked up as described for N-BOC-S-trityl-cysteine-4-aminobenzoyl methionine methyl ester in Example 1 and chromatographed on silica gel using a 1:1 hexanes and ethyl acetate elution mixture to give 0.12 g (56.0%) of pure material. ^1H NMR

(CDCl₃) 1.33 (9H, s), 1.61-1.68 (1H, m), 1.73-1.91 (4H, m), 1.96 (2H, t, J=7.6 Hz), 2.24 (6H, s), 2.57- 2.64 (2H, m), 3.57 (3H, s), 4.00 (1H, br s), 4.54-4.58 (1H, m), 5.84 (1H, d, J=7.8 Hz), 5.97 (1H, br d), 6.90 (1H, s), 6.92 (2H, s), 7.18-7.22 (9H, m), 7.27-7.40 (7H, m), 7.55 (1H, m), 7.61 (1H, m), 8.58 (1H, br s); ¹³C NMR (CDCl₃) 15.11, 21.20, 27.79, 29.25, 31.28, 51.70, 52.28, 54.08, 60.32, 71.45, 80.75, 118.01, 120.80, 126.38, 126.82, 127.98, 129.41, 129.87, 130.22, 138.11, 139.18, 139.79, 141.06, 144.17, 168.38, 169.04, 171.82.

G. TFA-Cysteine-4-amino-2-(3,5-dimethylphenyl)benzoyl methionine FTI275

15 N-BOC-S-trityl-cysteine-4-amino-2-(3,5-dimethylphenyl)benzoyl methionine methyl ester (0.12 g, 0.15 mmol) was placed into THF (0.9 ml) and was reacted with 0.5 M of LiOH (0.6 ml) at 0 °C as described above, followed by deprotection with
20 TFA (1.5 ml) and Et₃SiH (0.24 ml). Addition of excess scavenger does not appear to affect the result. The product was purified by preparative HPLC to give 23.8 mg (27.3%). mp 135-138°C; ¹H NMR (CDCl₃) 1.76-1.84 (1H, m), 2.00-2.17 (6H, m), 2.33 (6H, s), 3.05 (1H, dd, J=14.6 Hz, 7.3 Hz), 3.17 (1H, dd, J=14.6 Hz, J=4.9 Hz), 4.15 (1H, dd, J=7.3, 4.9 Hz), 4.45-4.48 (1H, m), 7.02 (3H, s), 7.53 (1H, d, J=8.0 Hz), 7.66 (2H, m); ¹³C (CD₃OD)
25 14.96, 21.51, 26.28, 30.91, 31.70, 53.03, 56.98, 119.27, 122.30, 127.52, 130.07, 130.57, 133.37, 139.28, 140.39, 141.29, 142.86, 166.89, 172.60, 174.81.

EXAMPLE 7

SYNTHESIS OF FTI-266**A. 4-amino-1-naphthoic acid**

4-amino-1-naphthalenecarbonitrile (1.50 g, 8.91 mmol) was dissolved into a 50% KOH solution (18 ml). The heterogeneous solution was heated at reflux for 2-3 days. Once the solution became homogenous and TLC showed no more starting material, the deep red solution was cooled and poured over 200 ml of water. The solution was then filtered and the acid was precipitated with concentrated HCl. The red crystals were filtered and the filtrate was refiltered to give pink crystals. The first fraction was treated with activated carbon to remove some of the red color. 1.51 g (90.6%) of product was obtained. mp 169-171°C; ¹H NMR (CD₃OD) 6.69 (1H, d, J=8.2 Hz), 7.38-7.43 (1H, m), 7.48-7.54 (1H, m), 8.03 (1H, d, J=8.5 Hz), 8.13 (1H, d, J=8.2 Hz), 9.09 (1H, d, J=8.5 Hz); ¹³C NMR (CD₃OD) 107.39, 114.61, 122.99, 123.92, 125.21, 127.40, 128.48, 135.04, 151.35, 171.44; HRMS calc. for C₁₁H₇NO₂, 187.0633; found, 187.0642.

B. N-BOC-4-amino-1-naphthoic acid

4-amino-1-naphthoic acid (0.86 g, 4.61 mmol) was dissolved into dioxane (9.2 ml) and 0.5 M NaOH (9.2 ml). Di-t-butyl dicarbonate (1.11 g, 5.07 mmol) was added and the mixture was stirred overnight. The reaction mixture was worked up as described for N-BOC-4-aminobenzoic acid in Example 1 to give 0.76 g (56.7%) of reddish pink solid. mp 194-195°C; ¹H NMR (CD₃OD) 1.56 (9H, s), 7.53-7.62 (2H, m), 7.79 (1H, d, J=8.1 Hz), 8.12 (1H, d, J=8.0 Hz), 8.22 (1H, d, J=8.18 Hz), 9.02 (1H, d, J=8.9 Hz); ¹³C NMR (CD₃OD), 26.68, 81.62, 119.06, 123.40, 124.57, 127.03, 127.37, 128.49, 128.77,

131.89, 133.76, 139.86, 155.95, 170.73; anal.
calc. for $C_{17}H_{17}NO_4$, C: 66.90, H: 5.96, N: 4.88;
found C: 66.49, H: 6.08, N: 4.79; m/z (EI), 289;
HRMS calc. for $C_{16}H_{17}NO_4$, 287.1158; found, 287.1151.

5 **C. N-BOC-4-amino-1-naphthoyl methionine methyl ester**

N-BOC-4-amino-1-naphthoic acid (0.46 g, 1.60 mmol), methionine methyl ester hydrochloride (0.35 g, 1.76 mmol), EDCI (0.43 g, 1.76 mmol), HOBT (0.24 g, 1.76 mmol) and Et_3N (0.27 ml) in CH_2Cl_2 (6.4 ml) were reacted as described for N-BOC-4-aminobenzoyl methionine methyl ester in Example 1. After workup and recrystallization from ethyl acetate and hexanes, 0.44 g (63.6%) of pale pink crystals were obtained. mp 131-132°C; 1H NMR ($CDCl_3$) 1.57 (9H, s), 2.11-2.21 (4H, m), 2.29-2.41 (1H, m), 2.65 (2H, t, $J=7.1$ Hz), 3.83 (3H, s), 4.99-5.06 (1H, m), 6.68 (1H, d, $J=8.0$), 7.02 (1H, s), 7.56-7.59 (2H, m), 7.69 (1H, d, $J=7.9$ Hz), 7.87-7.90 (1H, m), 8.02 (1H, d, $J=7.9$ Hz), 8.44-8.48 (1H, m); ^{13}C NMR ($CDCl_3$) 15.56, 28.31, 30.19, 31.65, 52.06, 52.64, 81.17, 115.82, 120.18, 125.79, 126.37, 126.53, 127.18, 131.02, 135.65, 152.93, 169.04, 172.40; HRMS calc. for $C_{22}H_{28}N_2O_5S$, 432.1719; found 432.1702; m/z (FAB) 433 (M+1).

25 **D. HCl·4-amino-1-naphthoyl methionine methyl ester**

N-BOC-4-amino-1-naphthoyl methionine methyl ester (0.57 g, 1.31 mmol) was deprotected with HCl/ ether to yield 0.31 g (64.1%) of white solid. mp 178-181°C; 1H NMR (CD_3OD) 2.08-2.16 (4H, m), 2.20-2.30 (1H, m), 2.57-2.75 (2H, m), 3.82 (3H, s), 4.87-4.91 (1H, m), 7.59 (1H, d, $J=7.5$ Hz), 7.67 (1H, d, $J=7.5$ Hz), 7.71-7.80 (2H, m), 8.03 (1H, dd, $J=7.1$ Hz, 2.0 Hz), 8.35 (1H, dd, $J=6.8$ Hz, 1.8 Hz); ^{13}C NMR (CD_3OD) 15.23, 31.40, 53.01,

53.33, 119.90, 122.20, 126.15, 127.41, 127.77,
129.09, 129.31, 131.50, 132.33, 135.64, 171.77,
173.83; m/z (FAB), 369 (M+1).

**E. N-BOC-S-trityl-cysteine-4-amino-1-naphthoyl
methionine methyl ester**

N-BOC-S-trityl-Cys (0.31 g, 0.67 mmol) in dry
THF (11.2 ml) was reacted with Et₃N (0.10 ml) and
IBCF (0.10 ml, 0.74 mmol) at -10 °C as described
above. HCl·4-amino-1-naphthoyl methionine methyl
ester (0.25 g, 0.67 mmol) in dry CH₂Cl₂ (3.5 ml)
was added and the mixture was stirred overnight
under nitrogen. The mixture was worked up as
described for N-BOC-S-trityl-cysteine-4-
aminobenzoyl methionine methyl ester in Example 1,
and the crude material was chromatographed on
silica gel using a 2:1 mixture of hexanes and
ethyl acetate to give 0.20g (37.5 %) of pure
product. ¹H NMR (CDCl₃) 1.48 (9H, s), 2.10-2.20
(4H, m), 2.30-2.37 (1H, m), 2.63 (2H, t, J=7.4),
2.74 (1H, J=12.9 Hz, J=5.3 Hz), 2.90 (1H, J=12.9
Hz, 6.2 Hz), 3.81 (3H, s), 4.96-5.03 (2H, m), 6.77
(1H, d, J=8.0 Hz), 7.18-7.33 (11H, m), 7.44-7.56
(7H, m), 7.60 (1H, d, J=7.7 Hz), 7.88 (1H, d,
J=8.0 Hz), 8.00 (1H, d, J=7.1 Hz), 8.37 (1H, d,
J=8.4 Hz), 8.94 (1H, br s); ¹³C NMR (CDCl₃) 15.23,
26.52, 31.41, 31.50, 52.98, 53.31, 56.79, 68.15,
122.52, 123.54, 126.16, 126.99, 128.03, 128.39,
129.52, 132.30, 134.04, 135.24, 168.08, 172.38,
173.94.

**F. TFA·cysteine-4-amino-1-naphthoyl methionine,
FTI-270**

N-BOC-S-trityl-cysteine-4-amino-1-naphthoyl
methionine methyl ester (83.3 mg, 0.11 mmol) was
taken up in THF (0.7 ml) and to this mixture was
added 0.5 M LiOH (0.43 ml) at 0°C. The mixture

was stirred at 0°C for 35 minutes. The solvent was removed in vacuo using a cold water bath. The residue was worked up as described for TFA·cysteine-4-amino-3-methylbenzoyl methionine in Example 2, and 74.1 mg of the free acid was obtained. This was then dissolved into CH₂Cl₂ (1 ml) and Et₃SiH (0.015 ml) was added followed by TFA (1 ml). The reaction mixture was stirred at room temperature for 1 h and worked up as further described for TFA·cysteine-4-amino-3-methylbenzoyl methionine in Example 2. After lyophilization, 42.4 mg of crude material was obtained which was then purified on the HPLC using 0.1% TFA in water and acetonitrile. mp 121-125°C; $[\alpha]_D^{25} = +2.4^\circ$ (c=0.8, H₂O); ¹H NMR (CD₃OD) 2.03-2.13 (4H, m), 2.22-2.36 (1H, m), 2.59-2.74 (2H, m), 3.16-3.33 (2H, m), 4.42 (1H, m), 4.84-4.89 (1H, m), 7.57-7.61 (2H, m), 7.64 (1H, d, J=7.7 Hz), 7.70 (1H, d, J=7.7 Hz), 8.08-8.11 (1H, m), 8.29-8.32 (1H, m), 8.98 (1H, d, J=7.7 Hz); ¹³C NMR (CD₃OD) 15.19, 26.45, 31.50, 31.63, 53.20, 56.72, 122.52, 123.43, 126.43, 126.12, 127.02, 127.96, 128.32, 129.49, 132.27, 134.15, 135.12, 168.11, 172.41, 175.17; anal. calc. for C₂₁H₂₃F₃N₃O₆S₂, C: 47.19, H: 4.34, N: 7.86; found, C: 46.53, H: 4.56, N: 7.59; Note: difference for C is 0.65.

G. HCl·cysteine-4-amino-1-naphthoyl methionine methyl ester FTI-270·HCl

TFA·cysteine-4-amino-1-naphthoyl methionine (0.12 g, 0.15 mmol) was dissolved in CH₃OH (4.3 ml). To this solution was added a solution of HgCl₂ (0.23 g, 0.86 mmol) in CH₃OH (4.3 ml). The procedure was continued as described above and after HCl/ Et₂O precipitation and several reprecipitations 31.0 mg (18.3 %) of pure white material was obtained. mp sub 137°C, decomp 214-

215°C; $[\alpha]_D^{25} = -32.0^\circ$ (c=1 CH₃OH); ¹H NMR (CD₃OD) 2.12 (3H, s), 2.21-2.28 (1H, m), 2.57-2.73 (3H, m), 3.20-3.34 (2H, m), 3.82 (3H, s), 4.39-4.43 (1H, m), 7.61-7.68 (3H, m), 7.78 (1H, d, J=7.7 Hz), 8.13-8.16 (1H, m), 8.28-8.32 (1H, m); ¹³C (CD₃OD) 15.23, 26.52, 31.41, 31.50, 52.98, 53.31, 56.79, 122.52, 123.54, 126.16, 126.99, 128.03, 128.39, 129.52, 132.30, 134.04, 135.24, 168.08, 172.38, 173.94.

10

EXAMPLE 8

SYNTHESIS OF FTI-254

A. N-Boc-S-trityl cysteinyl

Triethylamine (2.22 mL, 16 mmol) and N,O-dimethylhydroxylamine hydrochloride (1.57 g, 16.1 mmol) were added to a solution of N-Boc-S-trityl cysteine (7.44 g, 16 mmol) in 85 mL of methylene chloride. This mixture was cooled in an ice bath and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI, 3.08 g, 16.0 mmol) and HOBT (2.17 g, 16 mmol) was added. The mixture was stirred at 0°C for 1 hr and at room temperature for a further 10 hr. The mixture was extracted with methylene chloride and 0.5 N HCl. The organic layer was washed consecutively with 0.5 N HCl, concentrated NaHCO₃ and brine. The organic layer was dried and evaporated. The residue was purified by flash column chromatography (1.5 : 1 = hexane : ethylacetate) to give a white foam (7.40 g, 91%). m.p. 59-60 °C (decomp). ¹H NMR (CDCl₃) δ 7.41 (m, 6H), 7.20-7.31 (m, 9H), 5.13 (d, 8.9 Hz, 1H), 4.76 (br s, 1H), 3.64 (s, 3H), 3.15 (s, 3H), 2.56 (dd, 4.7 and 12.1 Hz, 1H), 2.39 (dd, 7.8 and 12.1 Hz, 1H), 1.43 (s, 9H). ¹³C NMR (CDCl₃) δ 170.7, 154.9, 144.2, 129.3, 127.6, 126.4, 79.3, 66.4, 61.2, 49.5, 33.8, 31.8, 28.1. This carboxyamide (2.02 g, 4.0 mmol) was dissolved in

30 mL of ether and cooled to -10 °C. Lithium aluminum hydride (167 mg, 4.40 mmol) was added and the mixture was stirred for 15 min under the nitrogen. Then 40 mL of 0.5 N HCl was added and the solution was extracted with ether. The ether layer was washed with 0.5 N HCl and dried. The evaporation of solvents gave a white foam (1.80 g) which was used for further reaction without purification. The ¹H NMR spectrum of this compound was complex. The percentage of the aldehyde was about 65-70%, which was calculated according to the integration of the sharp singlet (δ 9.17) and the trityl peak (δ 7.40, m, 6H; 7.28, m, 9H). Lowering the temperature to -45°C did not improve the aldehyde percentage.

B. 4-N-[2(R)-tert-Butoxycarbonylamino-3-triphenylmethylthiopropyl]aminobenzoyl methionine methyl ester.

One equivalent of N-Boc-S-trityl cysteinal in 10 mL of methanol was added to a solution of 4-aminobenzoyl methionine methyl ester hydrochloride (1.7836 g, 5.6 mmol) in 60 mL of methanol and 4 mL of glacial acetic acid. Sodium cyanoborohydride (0.528 g, 8.40 mmol) was added to this deep colored solution at 0 °C. The mixture was stirred at room temperature for 15 hr. After the evaporation of solvents, the residue was extracted with ethyl acetate and concentrated sodium bicarbonate. The organic phase was dried and the solvents were evaporated. The residue was purified through flash column chromatography (ethyl acetate / hexane = 1:1) to give a pure desired product (2.52 g, 65%). ¹H NMR (CDCl₃) δ 7.63 (d, 8.6 Hz, 2H), 7.43 (m, 6H), 7.21-7.32 (m, 9H), 6.73 (d, 7.6 Hz, 1H, Met amide), 6.50 (d, 8.6 Hz, 2H), 4.91 (ddd, 5.1 Hz, 5.3 Hz and 7.6 Hz, 1H, Met α H),

4.59 (d, 8.9 Hz, 1H, Boc amide), 4.25-4.40 (br, 1H, NHPh), 3.80 (m, 1H, Cys α H), 3.78 (s, 3H, OCH₃), 3.09 (d, 6.3 Hz, 2H, CH₂NH), 2.55-2.60 (m, 2H, CH₂SCPh₃), 2.46 (d, 5.0 Hz, 2H, CH₂SCH₃), 2.23-2.28 (m, 1H, Met CH₂), 2.07-2.12 (m, 1H, Met CH₂), 2.09 (s, 3H, SCH₃), 1.43 (s, 9H, Boc).

C. 4-N-[2(R)-Amino-3-mercaptopropyl]aminobenzoyl methionine methyl ester.

The fully protected 4-N-[2(R)-tert-Butoxycarbonylamino-3 triphenylmethylthiopropyl] amino-benzoyl methionine methyl ester (1.31 g, 1.83 mmol) was dissolved into 20 mL of methanol. To this solution was added mercuric chloride (1.09 g, 4.04 mmol) in 5 mL of methanol. The mixture was refluxed for 20 min and then cooled down. The clear solution was removed and the solid precipitate was washed with 5 mL of methanol. The solid was dried and then suspended in 15 mL of methanol. The suspension was stirred and reacted with hydrogen sulfide gas for 1 hr. The black precipitate was removed by centrifugation. The clear solution was evaporated to dryness. The residue was dissolved in 6 mL of methylene chloride followed by the addition of 20 mL of 3N HCl in ether. The white precipitate was filtered and dried to give a hydrochloride salt of the desired product (0.60 g, 73%). ¹H NMR (CD₃OD) δ 7.73 (d, 8.8 Hz, 2H), 6.75 (d, 8.8 Hz, 2H), 4.74 (dd, 4.9 Hz and 4.3 Hz, 1H, Met α H), 3.72 (s, 3H, OCH₃), 3.43-3.59 (m, 3H, CH₂NH and Cys α H), 2.93 (dd, 3.9 Hz and 14.4 Hz, 1H, CH₂SH), 2.81 (dd, 5.2 Hz and 14.5 Hz, 1H, CH₂SH), 2.49-2.66 (m, 2H, CH₂SCH₃), 2.07-2.20 (m, 2H, Met CH₂), 2.10 s, 3H, SCH₃).

D. 4-N-[2(R)-Amino-3-mercaptopropyl]aminobenzoyl methionine

The fully protected peptide 4-N-[2(R)-tert-Butoxycarbonylamino-3-triphenylmethylthiopropyl]-aminobenzoyl methionine methyl ester (567 mg, 0.79 mmol) was dissolved into 3.0 mL of 0.5 N lithium hydroxide and 3.0 mL of tetrahydrofuran. The mixture was stirred at 0°C for 1 hr. After the evaporation of solvents, the residue was dissolved in water and extracted with methylene chloride and 1N hydrochloric acid. The organic phase was dried and the solvents were evaporated. The residue was dissolved in a mixture of 1mL of methylene chloride and 2 mL of trifluoroacetic acid. Triethylsilane was added dropwise until the deep brown color disappeared. The mixture was kept at rt for 1 hr. The solvents were evaporated and the residue was dried. This residue was dissolved in 1 mL of 1.7N HCl in acetic acid followed by the addition of 20 mL of 3N HCl in ether. The white precipitate was filtered and dried to give a hydrochloride salt of the desired product (159 mg, 46%). Analytical HPLC showed purity over 98%. ¹H NMR (CD₃OD) δ 7.74 (d, 8.7 Hz, 2H), 6.75 (d, 8.7 Hz, 2H), 4.73 (dd, 4.5 Hz and 4.7 Hz, 1H, Met α H), 3.45-3.58 (m, 3H, CH₂NH and Cys α H), 2.93 (dd, 4.5 Hz and 14.6 Hz, 1 H, CH₂SH), 2.80 (dd, 5.3 Hz and 14.6 Hz, 1H, CH₂SH), 2.53-2.64 (m, 2H, CH₂SCH₃), 2.15-2.23 (m, 1H, Met CH₂), 2.07-2.13 (m, 1H, Met CH₂), 2.10 (s, 3H, SCH₃).

EXAMPLE 9

Synthesis of FTI-284

A. 4-Nitro-2-phenylbenzoyl-[1'(S)-methoxycarbonyl-3'-methylsulfonyl]propyl amide

4-nitro-2-phenylbenzoyl methionine methyl ester (525 mg, 1.28 mmol), N-methylmorpholine

oxide (453 mg, 3.87 mmol) and 0.5 mL of osmium tetroxide (2.5 wt.% solution in tert-butanol) were added to a mixture of 40 mL of acetone and 10 mL of water. The mixture was stirred at rt overnight. After the addition of excess sodium sulfite, the reaction mixture was extracted with ethyl acetate and washed with concentrated sodium bicarbonate. The organic phase was dried and the solvents were evaporated to give a solid (570 mg, 100%). ¹H NMR (CDCl₃) δ 8.29 (d, 7.7 Hz, 1H), 8.25 (s, 1H), 7.83 (d, 7.7 Hz, 1H), 7.43-7.55 (m, 5H), 6.15 (d, 7.3 Hz, 1H, Met amide), 4.68 (ddd, 5.0 Hz, 5.1 Hz and 7.3 Hz, 1H, Met α H), 3.70 (s, 3H, OCH₃), 2.85 (s, 3H, SCH₃), 2.69-2.81 (m, 1H, CH₂SO₂), 2.58-2.66 (m, 1H, CH₂SO₂), 2.21-2.33 (m, 1H, Met CH₂), 1.96-2.08 (m, 1H, Met CH₂).

B. 4-N-[2(R)-tert-Butoxycarbonylamino-3-triphenylmethylthiopropyl]amino-2-phenylbenzoyl-[1'(S)-methoxycarbonyl-3'-methylsulfonyl]propyl amide

The 4-Nitro-2-phenylbenzoyl-[1'(S)-methoxycarbonyl-3'-methylsulfonyl]propyl amide (430 mg, 1.02 mmol) was dissolved in 20 mL of methanol. A catalytic amount of 5% palladium on carbon was added and the mixture was hydrogenated at 40 PSI for 1.5 hr. The mixture was filtered and the filtrate was evaporated to dryness to give 4-amino product (400 mg, 100%). ¹H NMR (CD₃OD) δ 7.70 (d, 8.0 Hz, 1H), 7.38-7.47 (m, 7H), 4.53 (dd, 4.6 Hz and 4.8 Hz, 1H, Met α H), 3.72 (s, 3H, OCH₃), 2.89 (s, 3H, SO₂CH₃), 2.79-2.85 (m, 1H, CH₂SO₂), 2.58-2.68 (m, 1H, CH₂SO₂), 2.19-2.29 (m, 1H, Met CH₂), 1.93-2.04 (m, 1H, Met CH₂). This amine was dissolved in 15 mL of methanol and 0.8 mL of acetic acid. One equivalent of N-Boc-S-trityl cysteinal was added followed by the addition of sodium cyanoborohydride (97 mg, 1.5 eq). The

mixture was stirred at rt overnight. After the evaporation of solvents, the residue was extracted with ethyl acetate and concentrated sodium bicarbonate. The organic phase was dried and solvents were evaporated. The residue was purified through flash column chromatography (ethyl acetate / hexane / methanol = 15:15:2) to give a pure product (500 mg, 60%). ¹H NMR (CDCl₃) δ 7.64 (d, 8.5 Hz, 1H), 7.37-7.46 (m, 11H), 7.18-7.33 (m, 9H), 6.53 (d, 8.5 Hz, 1H), 6.34 (s, 1H), 5.74 (d, 7.5 Hz, 1H, Met amide), 4.64 (ddd, 4.9 Hz, 5.1 Hz and 7.5 Hz, 1H, Met α H), 4.55 (d, 7.5 Hz, 1H, Boc amide), 4.26 (br, 1H, NHPh), 3.79 (m, 1H, Cys α H), 3.68 (s, 3H, OCH₃), 3.10 (t, 5.7 Hz, 2H, CH₂NHPh), 2.84 (s, 3H, SO₂CH₃), 2.62-2.82 (m, 2H, CH₂SO₂), 2.45 (d, 2H, CH₂SCPh₃), 2.19-2.27 (m, 1H, Met CH₂), 1.84-1.95 (m, 1H, Met CH₂), 1.41 (s, 9H).

C. 4-N-[2(R)-Amino-3-mercaptopropyl]amino-2-phenylbenzoyl-[1'(S)-methoxycarbonyl-3'-methylsulfonyl]propyl amide, FTI-284

The fully protected peptide 4-N-[2(R)-tert-Butoxycarbonylamino-3-triphenylmethylthiopropyl]amino-2-phenylbenzoyl-[1'(S)-methoxycarbonyl-3'-methylsulfonyl]propyl amide (277 mg, 0.33 mmol) was dissolved into 5 mL of methanol. To this solution was added mercuric chloride (229 mg, 2.50 eq) in 2 mL of methanol. The mixture was refluxed for 20 min. The precipitate was dried and then suspended in 10 mL of methanol. This mixture was reacted with hydrogen sulfide gas. The reaction mixture was centrifuged and the clear solution was evaporated. The residue was dissolved in 2 mL of methylene chloride followed by the addition of 20 mL of 3N HCl in ether. The white precipitate was collected.

and dried to give a hydrochloride salt of the desired product (165 mg, 89%). ¹H NMR (CD₃OD) δ 7.44 (d, 8.4 Hz, 1H), 7.32-7.40 (m, 5H), 6.77 (d, 8.4 Hz, 1H), 6.68 (s, 1H), 4.45 (dd, 4.5 Hz and 4.7 Hz, 1H, Met α H), 3.69 (s, 3H, OCH₃), 3.40-3.57 (m, 3H, CH₂NHPh and Cys α H), 2.78-2.96 (m, 3H, CH₂SH and CH₂SO₂), 2.89 (s, 3H, SO₂CH₃), 2.60-2.69 (m, 1H, CH₂SO₂), 2.15-2.24 (m, 1H, Met CH₂), 1.91-2.02 (m, 1H, Met CH₂).

10

EXAMPLE 10

Synthesis of FTI-277

A. 4-N-[2(R)-tert-Butoxycarbonyl-3-triphenylmethylthiopropyl]amino-2-phenylbenzoyl methionine methyl ester

15

The coupling of 4-amino-2-phenylbenzoyl methionine methyl ester (3.88 g, 10 mmol) with one equivalent of N-Boc-S-trityl cysteinyl in the presence of 1.5 equivalent of sodium

20

cyanoborohydride gave a crude mixture which was purified through flash column chromatography

25

(ethyl acetate / hexane = 1:1) to give a pure desired product (5.83 g, 74%). ¹H NMR (CDCl₃) δ 7.65 (d, 8.5 Hz, 1H), 7.32-7.45 (m, 11H), 7.18-7.30 (m, 9H), 6.50 (d, 8.5 Hz, 1H), 6.33 (s, 1H), 5.65 (d, 7.6 Hz, 1H, Met amide), 4.62 (ddd, 5.0 Hz, 5.2 Hz and 7.6 Hz, 1H, Met α H), 4.54 (d, 8.1 Hz, Boc amide), 4.18 (br, 1H, NHPh), 3.78 (m, 1H, Cys α H), 3.64 (s, 3H, OCH₃), 3.10 (t, 6.1 Hz, 2H, CH₂NHPh), 2.45 (d, 5.0 Hz, 2H, CH₂SCPh₃), 2.04-2.10

30

(m, 2H, CH₂SCH₃), 2.00 (s, 3H, SCH₃), 1.81-1.92 (m, 1H, Met CH₂), 1.61-1.70 (m, 1H, Met CH₂), 1.40 (s, 9H). ¹³C NMR (CDCl₃) δ 172.0, 168.3, 155.7, 149.4, 144.3, 141.6, 141.1, 131.3, 129.5, 128.7, 128.5, 127.9, 127.7, 126.8, 122.6, 113.6, 111.3, 79.8, 67.1, 52.2, 51.7, 49.5, 47.2, 34.3, 31.6, 29.4, 28.3, 15.2.

35

B. 4-N-[2(R)-Amino-3-mercaptopropyl]amino-2-phenylbenzoyl methionine methyl ester, FTI-277.

The fully protected peptide 4-N-[2(R)-tert-Butoxycarbonyl-3-triphenylmethyl-thiopropyl]amino-2-phenylbenzoyl methionine methyl ester (1.57 g, 2.0 mmol) was first reacted with mercuric chloride (1.36 g, 5.0 mmol) and then reacted with hydrogen sulfide gas in methanol to give a hydrochloride salt of the desired product (0.808 g, 84%).

Analytical HPLC showed purity over 98%. $[\alpha]^{25}_D = -12.1^\circ$ ($c=0.008$, CH_3OH). ^1H NMR (CD_3OD) δ 7.42 (d, 8.3 Hz, 1H), 7.30-7.38 (m, 5H), 6.78 (d, 8.3 Hz, 1H), 6.71 (s, 1H), 4.47 (dd, 4.2 Hz and 5.1 Hz, 1H, Met α H), 3.68 (s, 3H, OCH_3), 3.44-3.54 (m, 3H, CH_2NHPh and Cys α H), 2.94 (dd, 4.1 Hz and 14.6 Hz, 1H, CH_2SH), 2.81 (dd, 5.0 Hz and 14.6 Hz, 1H, CH_2SH), 2.12-2.22 (m, 1H, CH_2SCH_3), 2.03-2.10 (m, 1H, CH_2SCH_3), 2.00 (s, 3H, SCH_3), 1.90-1.97 (m, 1H, Met CH_2), 1.73-1.82 (m, 1H, Met CH_2). ^{13}C NMR (CD_3OD) δ 173.7, 173.4, 150.7, 143.5, 142.3, 131.2, 129.8, 129.5, 128.6, 125.6, 115.6, 112.2, 53.7, 53.2, 52.8, 45.0, 31.3, 30.8, 25.3, 15.0.

EXAMPLE 11

Synthesis of FTI-276

4-N-[2(R)-Amino-3-mercaptopropyl] amino-2-phenylbenzoyl methionine

The fully protected peptide 4-N-[2(R)-tert-Butoxycarbonyl-3-triphenylmethylthiopropyl]-amino-2-phenylbenzoyl methionine methyl ester (2.36 g, 3 mmol) was first reacted with lithium hydroxide and then with trifluoroacetic acid to give a crude product (1.30 g, 77% yield, 85% purity shown by HPLC) which was further purified through preparative HPLC to give a pure product (0.98 g, 75%). $[\alpha]^{25}_D = -13.6^\circ$ ($c=0.005$, CH_3OH). ^1H NMR (CD_3OD) δ 7.44 (d, 8.4 Hz, 1H), 7.30-7.41 (m, 5H),

6.75 (d, 8.4 Hz, 1H), 6.68 (s, 1H), 4.43 (dd, 4.2 Hz and 5.1 Hz, 1H, Met α H), 3.44-3.58 (m, 3H, CH₂NHPh and Cys α H), 2.95 (dd, 4.4 Hz and 14.5 Hz, 1H, CH₂SH), 2.83 (dd, 5.0 Hz and 14.5 Hz, 1H, CH₂SH), 2.14-2.23 (m, 1H, CH₂SCH₃), 2.05-2.11 (m, 1H, CH₂SCH₃), 2.00 (s, 3H, SCH₃), 1.91-1.99 (m, 1H, Met CH₂), 1.72-1.81 (m, 1H, Met CH₂). ¹³C NMR (CD₃OD) δ 176.4, 173.5, 150.4, 143.0, 141.5, 131.0, 129.7, 129.4, 128.9, 124.6, 115.0, 112.2, 53.3, 44.4, 30.8, 30.1, 24.9, 14.8.

Other compounds of the invention (in particular those of claims [[14-18]) are synthesizable by modifications of the procedure described for the 2-phenyl-4-aminobenzoic acid derivative of claim 3. In particular, modifications of the Suzuki coupling method will allow the incorporation of an alkoxy-, chloro, bromo or methyl substituted phenyl group onto the 4-aminobenzoic acid spacer. As with the unsubstituted derivative, 4-nitro-2-bromotoluene will be coupled with the corresponding substituted phenyl boronic acid derivative (alkoxyphenyl or chloro-, bromo- or methylphenylboronic acid) under palladium catalyzed conditions. The appropriately substituted 2-(substituted) phenyl-4-nitro toluene derivative will be incorporated into the peptidomimetic synthesis as described for the 2-phenyl case.

In a similar way the precursor to the 2-naphthyl-, 2-thiophene-, 2-pyrrole-, and 2-pyridyl-4-aminobenzoic acid spacers can be prepared by reaction of 4-nitro-2-bromotoluene with naphthalene-2-boronic acid, thiophene-2-boronic acid, pyrrole-2-boronic acid, pyridine-2,3- or 4-boronic acid.

EXAMPLE 12

FTase and GGTase I Activity Assay

FTase and GGTase I activities from 60,000 X g supernatants of human Burkitt lymphoma (Daudi) cells (ATCC, Rockville, MD, USA) were assayed as described previously for FTase (41). Briefly, 100 μ g of the supernatant was incubated in 50 mM Tris, pH 7.5, 50 μ M ZnCl₂, 20 mM KCl and 1 mM dithiothreitol (DTT). The reaction was incubated at 30°C for 30 min with recombinant Ha-Ras-CVLS (11 μ M) and [³H]FPP (625 nM; 16.3 Ci/mmol) for FTase, and recombinant Ha-Ras-CVLL (5 μ M) and [³H]geranylgeranylpyrophosphate (525 nM; 19.0 Ci/mmol) for GGTase I. The peptidomimetics were mixed with FTase and GGTase before adding to the reaction mixture.

EXAMPLE 13

Ras and Rap1A Processing Assay

H-RasF cells (45) were seeded on day 0 in 100 mm Dishes (costar) in Dulbecco's modified Eagles medium (GIBCO) and allowed to grow to 40-60% confluency. On days 1 and 2, cells were fed with 4 ml of medium per plate plus various concentrations of FTI-277 or vehicle. On day 3, cells were washed one time with ice cold PBS and were collected and lysed by incubation for 30-60 min on ice in lysis buffer (41). Lysates were cleared (14,000 rpm, 4°C, 15 min) and supernatants collected. Equal amounts of lysate were separated on a 12.5% SDS-PAGE, transferred to nitrocellulose, and a western blot performed using a anti-Ras antibody (Y13-238, ATCC) or anti-Rap1A antibody (Santa Cruz Biotechnology, Santa Cruz, CA). Antibody reactions were visualized using peroxidase-conjugated goat anti-rat IgG for Y13-238 and peroxidase-conjugated goat anti-rabbit IgG

for Rap1A and an enhanced chemiluminescence detection system (ECL; Amersham Corp.)

EXAMPLE 14

Co-immunoprecipitation of Raf and Ras

5 Cells were seeded on day 0 in 100 mm dishes
in 10 ml Dulbecco's Modified Eagles Medium
(GIBCO) supplemented with 10% calf serum (Hyclone)
and 1% pen/strep (GIBCO). On days 1 and 2 cells
10 were treated with FTI-277 (5 μ M) or vehicle
(confluency of cells 40-60%). On day 3, cells
were collected by centrifugation in ice cold PBS.
Cell pellets were then resuspended in ice cold
hypotonic buffer (10 mM Tris, pH 7.5, 5 mM MgCl₂, 1
mM DTT, 1 mM PMSF) and cells were sonicated to
15 break up cell pellet to promote separation of
cytosol and membrane. The cell suspension was
then centrifuged at 2,000 rpm for 10 min to clear
debris after which the supernatant was loaded in
ultracentrifuge tubes and spun for 30 min at
20 100,000 X g to SW Ti55 Rotor to separate membrane
and cytosol fractions. The cytosol and membrane
fractions were lysed on ice for 60 min in buffer
containing 30 mM HEPES, pH 7.5, 1% TX-100, 10%
glycerol, 10 mM NaCl, 5 mM MgCl₂, 2 mM Na₃VO₄, 25
25 mM NaF, 1 mM EGTA, 10 μ M soybean trypsin
inhibitor, 25 μ g/ml leupeptin, 10 μ g/ml aprotinin,
2 mM PMSF). The lysates were clarified by
centrifugation. Equal amounts of cytosol and
membrane fractions were immunoprecipitated using
30 50 μ l of a 25% Protein-A Sepharose Cl-4B
suspension (Sigma) with 1 μ g/ml anti-c-Raf-1
(SC133, Santa Cruz Biotechnology, Santa Cruz, CA).
The samples were tumbled at 4°C for 60 min and then
washed 5 times in 50 mM HEPES, pH 7.5, 100 mM
35 NaCl, 5 mM MgCl₂, 0.1% TX-100, 10% glycerol, 20 mM
NaF. The final pellets were run on 12.5% SDS-

PAGE, transferred to nitrocellulose, and immunoblotted for the presence of Ras using anti-Ras antibody (Y13-238) and immunoblotted for the presence of Raf (c-Raf-1, SC133, Santa Cruz Biotechnology, Santa Cruz, CA). Detection was the same as above for Ras and Rap1A processing.

EXAMPLE 15

Detection of GTP and GDP bound to Ras (FTI-277)

H-RasF cells were seeded and treated as above for Ras/Raf interaction and Ras and Rap1A processing. On day 2, however, cells were labeled overnight with [³²P] orthophosphate at 100 μ Ci/mo (Amersham PBS13) in 10 ml DMEM-phosphate supplemented with 10% calf serum, 1 mg/ml BSA and 20 mM HEPES, pH 7.5. On day 3, the medium was removed and cells were washed one time in ice-cold PBS, scraped from the plate with a cell scraper, collected and centrifuged. The cell pellet was resuspended in ice-cold hypotonic buffer listed above and the cytosol and membrane fractions were separated according to the above description for Ra/Raf association. The cytosol and membrane fractions were lysed on ice for 60 min in 50 mM Tris, pH 7.5, 5 mM MgCl₂, 1% Triton X-100 (TX-100), 0.5% DOC, 0.05% SDS, 500 mM NaCl, 1 mM EGTA, 10 μ g/ml aprotinin, 10 μ g/ml soybean trypsin inhibitor, 25 μ g/ml leupeptin, 1 mM DTT, 1 mg/ml BSA. Lysates were cleared and equal amounts of protein were immunoprecipitated using anti-Ras antibody (Y13-259) along with 30 μ l Protein A-Agarose goat anti-rat IgG complex (Oncogene Science) for 60 min at 4°C. Immunoprecipitates were washed 6 times in 50 mM HEPES, pH 7.5, 0.5 M NaCl, 0.1% TX-100, 0.0005 SDS, 5 mM MgCl₂, drained using a syringe and bound nucleotide eluted in 12 μ l of 5 mM DTT, 5 mM EDTA, 0.2% SDS, 0.5 mM GDP

and 0.5 mM GTP at 68°C for 20 min. Immune complexes were spun down quickly and 6 μ l of the supernatant was loaded onto PEI cellulose thin layer chromatography plates (20 cm X 20 cm).
5 Nucleotides were separated by chromatography in 78 g/linter ammonium formate, 9.6% (v/v) concentrated HCl. Plates were analyzed by autoradiogram.

EXAMPLE 16

Analysis of Raf-I Kinase Activity

10 Raf-1 kinase was assayed by determining the ability of Raf to transfer phosphate from [γ - 32 P] ATP to a 19-mer peptide containing an autophosphorylation site. Membrane and cytosol fraction isolation and Raf immunoprecipitates were
15 washed three times with cold HEPES buffer and twice with kinase buffer (50 mM Tris, pH 7.5, 150 mM NaCl, 12 mM MnCl₂, 1 mM DTT, 0.2% Tween 20. Immune complex kinase assays were performed by incubating immunoprecipitates from membrane and
20 cytosol fractions in 96 μ l of kinase buffer with 20 μ Ci of [γ - 32 P]ATP (10 mCi/ml, Amersham) and 2 μ l of the Raf-1 substrate peptide (1 mg/ml, Promega) for 30 min at 25°C. The sequence of the Raf-1 substrate peptide is IVQQFGFQRRASDDGKLTD.
25 The phosphorylation reaction was terminated by spotting 50 μ l aliquots of the assay mixture onto Whatman P81 for 40 min in 0.5% orthophosphoric acid and air dried. The amount of 32 P incorporated was determined by liquid scintillation counting.

30

EXAMPLE 17

Inhibition of FTase by FTI-276 and other compounds

Fig. 1B shows that FTI-276 inhibited the transfer of farnesyl from [3 H]FPP to recombinant H-Ras-CVLS with an IC₅₀ of 500 pM. FTI-249, the
35 parent compound of FTI-276, inhibited FTase with

an IC_{50} of 200,000 pM. Thus, a phenyl ring at the 2 position of the benzoic acid spacer increased inhibition potency of FTase by 400 fold confirming our prediction of a significant hydrophobic pocket within the CAAX binding site of FTase. This extremely potent inhibitor was also highly selective (100-fold) for FTase over the closely related GGTase I (Fig. 1B). FTI-276 inhibited the transfer of geranylgeranyl from [3H]GG-PP to recombinant H-Ras-CV11 with an IC_{50} of 50 nM (Fig. 1B). This 100-fold selectivity is superior to the 15-fold selectivity of the parent compound, FTI-249. Data for a number of other compounds of interest are shown in Table 1.

Table 1

Compound		FTase IC ₅₀ (nM)	GGTase I IC ₅₀ (nM)	GG/F
FTI				
232	CABAM	213	1200	6
260	3-Me-CABAM	825	9000	11
261	3-OMe-CABAM	2550	50000	20
270	CANAM	143	3150	22
272	2-Ph-CABAM	5	267	53
274	2-Ph-CABAM-OMe	2050	30000	15
275	2-Xy-CABAM	405	400	1
249	red.CABAM	272	3967	15
254	red.CABAM-OMe	1000	19000	19
276	red.2-Ph-CABAM	0.5	57	114
277	red.2-Ph-CABAM-OMe	50	1600	32

EXAMPLE 18

Inhibition of Ras Processing by FTI-277

To facilitate cellular uptake, FTI-277, the methylester of FTI-276, was used in experiments to measure inhibition of Ras processing. H-RasF cells (NIH 3T3 cells transformed with oncogenic (61 leucine) H-Ras-CVLS (45) were treated with FTI-277 (0-50 μ M) and the lysates blotted with anti-Ras or anti-Rap1A antibodies. As shown in Fig. 2A, concentrations as low as 10 nM inhibited Ras processing but concentrations as high as 10 μ M did not inhibit processing of the geranylgeranylated Rap1A. FTI-277 inhibited Ras processing with an IC₅₀ of 100 nM. In contrast, the IC₅₀ of FTI-249 is 100 μ M, and the most potent CAAX peptidomimetics previously reported inhibited

Ras processing at concentrations of 10 μ M or higher (44).

The selectivity of FTI-277 for farnesylation but not geranylgeranylation processing is further demonstrated in Fig 2B. H-RasGG cells (NIH 3T3 cells transformed with oncogenic (61 leucine) H-Ras-CVLL (45) were treated with FTI-277.

Processing of RasGG was not affected, whereas that of RasF was completely blocked. The processing of endogenous Ras is also blocked in pZIPneo cells (NIH 3T3 cells transfected with the same plasmid as H-RasF and H Ras FF except the vector contained no oncogenic Ras sequences) and Raf cells (NIH 3T3 cells transformed by an activated viral Raf (48)).

Mechanism of disruption of Ras oncogenic signalling by FTI-277

Ras relays biological information from tyrosine kinase receptors to the nucleus by activation of a cascade of MAPKs (reviewed in 29-31). Upon growth factor stimulation, Ras becomes GTP bound and is then able to recruit the ser/thr kinase c-Raf-1 to the plasma membrane where it is activated. c-Raf-1 then phosphorylates and activates MEK, a dual thr/tyr kinase, which activates MAPK. Recently, epidermal growth factor has been shown to induce association of Raf with Ras (46).

In order to determine the mechanism by which FTI-277 disrupts Ras oncogenic signaling, NIH 3T3 cells were transfected with activated (GTP-locked) Ras and the effects of FTI-277 on the interaction of Ras with its immediate effector, Raf, were investigated. Various NIH 3T3 cell transfectants (pZIPneo, H-RasF, and H-RasGG) were treated with vehicle or FTI-277, membrane and cytosolic fractions were isolated and immunoprecipitated

with anti-Raf antibody as described above. Raf did not associate with Ras in pZIPneo cells which did not contain GTP-locked Ras, as shown in Fig. 3. In contrast, H-RasF and H-RasGG cells contain Ras/Raf complexes in the membrane, but not in the cytosolic fractions, as shown in Fig. 3. Treatment of these cells with FTI-277 resulted in the accumulation of Ras/Raf complexes in the cytosolic but not membrane fractions of H-RasF cells, but not in the H-RasGG cells (Fig 3). Thus, the disruption of Ras/Raf interaction at the cell membrane and accumulation of these complexes in the cytoplasm occurred only in Ras-F but not Ras-GG cells, in agreement with the Ras processing selectivity results of Fig. 2. Thus, these results demonstrate that inhibition with FTI-277 results in the accumulation of non-farnesylated cytosolic Ras that is capable of binding to Raf. The fact that non-processed Ras can associate with Raf in a non-membranous cytoplasmic environment was confirmed by transfecting NIH 3T3 cells with a GTP-locked Ras that lacks a farnesylated site and, therefore, remains in the cytoplasm (Ras mutant with a 61 leucine oncogenic mutation and a 186 serine mutation) and showing that these cells contained only cytoplasmic Ras/Raf complexes when immunoprecipitated with Raf and blotted with antiRas antibodies (Fig. 3). In short, farnesylation is not required for Ras to bind to Raf.

EXAMPLE 19

Determination of nucleotide state of Ras

The fact that Raf binds Ras-GTP with much higher affinity than Ras-GDP was used to determine the nucleotide state of Ras in the cytoplasmic Ras/Raf complexes, as described above. In Ras-F

cells, only membrane fractions contained GTP-locked Ras, as shown in Fig 4A. Upon treatment with FTI-277, however, the non-farnesylated cytosolic Ras was found to be GTP bound. Thus, binding of GTP to 61 leucine Ras does not require Ras processing and subsequent plasma membrane association. The ser/thr kinase activity of Raf in Ras/Raf complexes was then determined by immunoprecipitating Raf and assaying for its ability to phosphorylate a 19-mer autophosphorylated peptide. Fig. 4B shows that oncogenic Ras-F induced activation of Raf in the plasma membrane and that treatment with FTI-277 suppressed this activation. More importantly, the cytoplasmic Ras/Raf complexes that were induced by FTI-277 (Fig. 3) had basal levels of Raf kinase activity that were comparable to those of the parental NIH 3T3 cell line pZIPneo (Fig. 4B). Taken together, Figures 3 and 4 demonstrate that oncogenic transformation with GTP-locked Ras results in the constitutive recruitment to the plasma membrane and subsequent activation of Raf. Furthermore, FTase inhibition by FTI-277 suppresses this activation by inducing the accumulation of Ras/Raf complexes in the cytoplasm where Ras is GTP-bound but Raf kinase is not activated. The fact that Raf kinase is not activated when bound to Ras in a non-membranous environment is consistent with recent reports that indicate that Raf activation requires an as yet to be determined activating factor at the plasma membrane (47).

Experiments were then performed to investigate the effects of FTI-277 on oncogenic Ras activation of MAPK, a Raf downstream signalling event (29-31). Oncogenic activation of MAPK can be easily detected since activated MAPK

migrates slower in SDS-PAGE. Fig. 5A shows that NIH 3T3 cells transfected with pZIPneo contain only inactive MAPK but that upon transformation with oncogenic H-Ras, MAPK is activated (Fig. 5A).

5 Pretreatment with FTI-277 results in a concentration dependent inhibition of oncogenic Ras activation of MAPK. Concentrations as low as 300 nM were effective and the block was complete at 1 μ M. Taken together, Figs. 3 and 5
10 demonstrate that at least 50% inhibition of Ras processing is required for complete suppression of MAPK activation but that less than a 100% inhibition of Ras processing is required for complete suppression of MAPK activation by Ras. A
15 series of NIH 3T3 cell lines transformed with various oncogenes was used to determine whether the inhibition of MAPK activation is due to selectively antagonizing Ras function. Fig. 5B shows that FTI-277 was able to block H-RasF but
20 not H-RasGG activation of MAPK. This is consistent with its ability to inhibit H-RasF but not H-RasGG processing (Fig. 2). Selectivity of FTI-277 towards antagonizing Ras-dependent
25 activation of MAPK was substantiated by using NIH 3T3 cells where MAPK is constitutively activated by transformation with the Raf oncogene. Fig. 5B shows that oncogenic Raf activation of MAPK is not blocked by FTI-277 even though processing of endogenous Ras was inhibited in these cells.
30 Similar results were also obtained with FTI-276 (Fig. 6). Taken together these results clearly demonstrate that FTI-276 and FTI-277 are highly effective and selective in disrupting constitutive Ras-specific activation of MAPK.

35 Thus, FTI-277 is an extremely potent and highly selective FTase inhibitor. This compound inhibited Ras processing with concentrations as

low as 10 nM and processing was blocked at 1 μ M. The most potent inhibitor previously reported BZA-5B, blocked Ras processing only at 150 μ M (44).

EXAMPLE 20

5 Antitumor Efficacy and Selectivity of FTI-276 and
 FTI-277 In order to demonstrate the efficacy of
 these inhibitors as anticancer agents and show
 that they can inhibit tumor growth of human tumors
 which have multiple and complex genetic
10 alterations, antitumor efficacy experiments were
 performed using a human tumor cell line. A
 critical issue connected with the potential use of
 the compounds of the invention is whether the
 growth of human tumors which harbor K-Ras
15 mutations can be blocked. This is important for
 further development of FTase inhibitors as
 anticancer drugs since K-Ras mutations are most
 common in human cancers and since K-Ras processing
 is more difficult to inhibit than the processing
20 of the less prevalent H-Ras (1-3, 15).
 Furthermore, the majority of human tumors have
 multiple genetic alternations; notably a deletion
 in the tumor suppressor gene p53 is most
 prevalent. It is therefore extremely important to
25 determine whether or not inhibition of Ras
 function is sufficient to halt the growth of human
 tumors which harbor K-Ras mutation as well as
 deletions in p53.

 To evaluate the antitumor efficacy of FTI-
30 276, a nude mouse xenograft model was used. In
 this model, tumors from two human lung carcinoma
 cell lines are implanted subcutaneously. One of
 these (Calu-1) harbors a K-Ras oncogenic mutation
 and has a deletion of the tumor suppressor gene
35 p53. The other human lung carcinoma (NCI-H180)
 has no Ras mutations. Thirty two days after s.c.

implantation when the tumors reached sizes of 60 to 80 mm³, the mice were randomly separated into control and treated groups (4 animals per group; each animal had a tumor on both the right and the left flank). Figure 7A shows that tumors from control animals treated with saline once daily starting on day 36 grew to an average size of 566 mm³ over a period of 64 days from tumor implantation. In contrast, tumors treated once daily with FTI-276 (50 mg/kg) grew very little and the average tumor size was 113 mm³ (Fig. 7A). In another experiment, FTI-277, the methylester of FTI-276, inhibited the growth of Calu-I cells to the same extent (Figure 8). Although the animals were treated once daily with 50 mg/kg for 36 days (total cumulative does of 1.8 g/kg), no weight loss was observed and the animals appeared normal with no evidence of gross toxicity. This lack of toxicity was also observed in separate experiments where the dose was escalated to 180 mg/kg once daily. Thus, FTI-276 and FTI-277 essentially blocked tumor growth of Calu-I carcinoma with no evidence of gross toxicity.

The effect of FTI-276 on the tumor growth of another human lung carcinoma, NCI-H810, that does not harbor an oncogenic Ras mutation was also determined. Figure 7B shows that tumors from animals treated with saline or FTI-276 grew at a similar rate. Over a period of 14 days of treatment the average tumor size of the control and FTI-276 treated groups were 919 mm³ and 815 mm³, respectively. These results clearly demonstrate that in contrast to Calu-1, NCI-H810 carcinomas were not sensitive to FTI-276 treatment suggesting that FTI-276 inhibition of tumor growth of human lung carcinomas is Ras-dependent.

Furthermore, FTI-276 inhibited tumor growth even though Calu-1 does not express p53.

To further establish the selectivity of FTI-276 to inhibit selectively Ras-dependent tumors, the anti-tumor efficacy of FTI-276 and FTI-277 against H-RasF and Raf transformed NIH 3T3 in the same nude mouse xenograft model was examined. Figure 9 shows that a once daily injection of FTI-276 or FTI-277 (50 mg/kg) inhibited tumor growth of H-RasF transformed NIH 3T3 cells. In contrast, an identical treatment regimen with FTI-276 and FTI-277 had no effect on the growth of Raf-transformed NIH 3T3 cells (Fig. 10), further confirming the conclusion from the results of Figures 7 and 8 that FTI-276 and FTI-277 are selective for Ras-dependent tumors.

In addition, the question of whether FTI-276 inhibition of tumor growth correlated with inhibition of Ras processing *in vivo* was addressed. To do this, mice having subcutaneous H-RasF cells were treated with various doses of FTI-276 (0, 10, 50 and 100 mg/kg) and tumor size and Ras processing in the HRasF tumors *in vivo* were examined. Figure 11A shows that throughout the 11 day treatment period, FTI-276 inhibited tumor growth in a dose dependent fashion. The tumor sizes at the end of 17 days were 2490 mm³ for saline, 1793 mm³ for 10 mg/kg, 1226 mm³ for 50 mg/kg and 624 mm³ for 100 mg/kg treated animals. To determine the levels of inhibition of Ras processing, the animals were sacrificed 5 hrs after the last injection, the tumors were excised and processed for immunoblotting with anti-Ras antibody as described in legend to Fig. 11. Tumors from control animals contained only fully processed Ras which migrates faster in SDS-PAGE gels (Fig. 11B). As the dose of FTI-276 increases

from 10 to 100 mg/kg there was a progressive accumulation of unprocessed Ras which was paralleled by a decrease in the relative ratio of fully processed Ras. Thus, the extent of tumor growth inhibition correlated with the extent of inhibition of Ras processing. Furthermore, the inhibition of Ras processing in vivo was selective in that FTI-276 did not inhibit Rap1A processing even at 100 mg/kg.

II. Farnesyltransferase Inhibitors of the type CA

Compounds of another major embodiment of the invention are represented by formula II. Several examples are shown in Figure 12. These and other compounds of this embodiment may be prepared using procedures which are conventional in the art. For example, compounds 4 and 5 of Figure 12 may be prepared by reductive amination of 4-amino-3'-tert.butoxy-carbonyl biphenyl or 4-amino-4'-tert.butoxy carbonyl biphenyl, respectively, with N-Boc-S-trityl cysteinal followed by deprotection with, for example, trifluoroacetic acid and purification.

This embodiment of the invention is illustrated but not limited by the following examples:


EXAMPLE 21

The compound C-4ABA-Met of formula (2) (see Figure 12) was prepared as described in reference (27). The protected form of the peptidomimetic (2a) was prepared through the reductive amination of 4-aminobenzoyl methionine methyl ester and N-Boc-S-trityl cysteinal in methanol solution containing NaBH_3CN and 5% acetic acid. This reaction gave the N-Boc-S-trityl, methyl ester of (2a) with a yield of 65%. The protected


peptidomimetic was deesterified by LiOH in THF and then deprotected by trifluoroacetic acid in methylene chloride with two equivalents of triethylsilane to give crude (2a) which was purified by reverse phase HPLC. The biphenyl-based peptidomimetic (8) was prepared by the reductive amination of 4-amino-3'-methyl biphenyl with N-Boc-S-trityl cysteinal, to give the N-Boc-S-trityl derivatives of (8), which was then deprotected by trifluoroacetic acid and purified by reverse phase HPLC. The peptidomimetics (4) and (5) were prepared from the reductive amination of 4-amino-3'-tert.butoxycarbonyl biphenyl and 4-amino-4'-tert.butoxycarbonylbiphenyl, respectively, with N-Boc-S-trityl cysteinal, to give the N-Boc-S-trityl, tert-butyl ester of (4) and (5). Deprotection by trifluoroacetic acid and purification by reverse phase HPLC gave pure (4) and (5).

Synthesis

The basic approach used for the preparation of the compounds of the invention is illustrated in Scheme 1 with the synthesis of compound 4. [Compound numbers in the following discussion refer to Schemes 1 and 2.]

Scheme 1. Representative Synthesis of FTase Inhibitors^a

^a Reagents: (a) Pd(OAc)₂; (b) KMnO₄, pyridine/H₂O; (c) (1) (COCl)₂, (2) *tert*-butyl alcohol, *n*-BuLi; (d) (1) H₂, Pd/C, (2) *N*-Boc-S-tritylcysteine 17, (3) NaB(CN)H₃; (e) TFA, Et₃SiH.

Scheme 2. Synthesis of Compounds 11 and 12^a

1-Bromo-4-nitrobenzene was coupled to 3-methylphenylboronic acid (34) through an modified Suzuki coupling (30) to afford compound 14 (35). Compound 14 was oxidized to carboxylic acid 15 which was converted to the acid chloride and reacted with lithium tert-butoxide (36) to give the tert-butyl ester 16. Reduction of 16 by hydrogenation and subsequent reductive amination (37) of the resulting amine with N-Boc-S-trityl-cysteinyl 17 (38) gave the fully protected derivative 18, which was deprotected by trifluoroacetic acid in the presence of triethylsilane (39). Compound 4 was purified by reverse phase HPLC and isolated as its trifluoroacetate salt by lyophilization.

The synthesis of 11 and 12 is described in Scheme 2. Compound 19 was made from 3-methyl-3'-carboxybiphenyl (itself formed from aryl-aryl coupling of methyl 3-bromobenzoate with 3-methylphenylboronic acid followed by a saponification) via the same method as compound 16. Bromination of 19 followed by reaction with sodium azide gave 20 which was catalytically hydrogenated to give the corresponding amine. Reaction of Boc-trityl protected cysteine with this amine through the mixed anhydride method gave 21, while compound 22 was made by reductive amination with Boc-trityl protected cysteinyl.

Experimental

¹H and ¹³C NMR spectrum were recorded on a Bruker AM-300 spectrometer. Chemical shifts were reported in δ (ppm) relative to tetramethylsilane. All coupling constants were described in Hz. Elemental analyses were performed by Atlantic Microlab Inc., Georgia. Optical rotations were measured on a Perkin-Elmer 241 polarimeter.

Concentrations are expressed in g/mL. Flash column chromatography was performed on silica gel (40-63 μ m) under a pressure about 4 psi. Solvents were obtained from commercial suppliers and purified as following: tetrahydrofuran and ether were distilled from sodium benzophenone ketyl, methylene chloride was distilled over lithium aluminum hydride. Preparative HPLC was performed using a Waters 600 E controller and a Waters 490 E Multi-Wavelength UV detector with a 25x10 cm Delta-Pak C-18 300 Å cartridge column inside a Waters 25x10 cm Radial Compression Module. Analytical HPLC was performed using a Rainin HP Controller and a Rainin UV-C detector with a Rainin 250x4.6 mm 5 μ m Microsorb C-18 column. High resolution mass spectra (HRMS) and low resolution mass spectra (LRMS) were performed on a Varian MAT CH-5 and VG 7070 mass spectrometer. The purity of all the synthesized inhibitors was more than 98% as indicated by analytical HPLC.

A. 4-Nitro-3'-methylbiphenyl (14).

To a mixture of 4-nitrobenzene (3.0 g, 14.8 mmol) and 3-methylphenylboronic acid (2.06 g, 15.1 mmol) in 35 mL of acetone and 40 mL of water was added $K_2CO_3 \cdot 1.5H_2O$ (5.93 g, 37.5 mmol) and $Pd(OAc)_2$ (101 mg, 0.50 mmol). The deep black mixture was refluxed for 6 hr and then cooled. The mixture was extracted with ether and the organic layer was passed through a layer of celite. The pale yellow solution was dried over Na_2SO_4 and evaporated to dryness. The residue was recrystallized from hot methanol to give pale yellow crystals (2.68 g, 85%). m.p. 59-60 °C. 1H NMR ($CDCl_3$) δ 8.26 (d, 8.7 Hz, 2H), 7.70 (d, 8.7 Hz, 2H), 7.41 (m, 3H), 7.26 (d, 7.1 Hz, 1H), 2.43 (s, 1H). ^{13}C NMR ($CDCl_3$) δ 147.6, 146.8, 138.8, 138.6, 129.6, 128.9, 128.0,

127.6, 124.4, 123.9, 21.4. LRMS (EI) for $C_{13}H_{11}NO_2$ 213 (M^+ , intensity 100); HRMS (EI) calcd 213.0789, obsd 213.0778.

B. 4-Nitro-3'-carboxybiphenyl (15).

5 Compound 14 (2.31 g, 10 mmol) was suspended in a mixture of 10 mL of pyridine and 20 mL of water. The mixture was heated to refluxing and then $KMnO_4$ (7.9 g, 50 mmol) was added in portions. This mixture was refluxed for 1 hr and then
10 stirred at room temperature for 4 hr. The hot mixture was filtered and the black solid was washed with hot water. The filtrate was acidified with 6 N HCl. The precipitate was collected and dried (2.16 g, 89%). m.p. 265°C (decomp). 1H NMR
15 (DMSO- d_6) δ 11.1-11.4 (br s, COOH), 8.32 (d, 8.7 Hz, 2H), 8.27 (s, 1H), 8.02 (m, 4H), 7.66 (t, 7.8 Hz, 1H). ^{13}C NMR (DMSO- d_6) δ 167.1, 148.9, 145.6, 138.2, 131.8, 131.5, 129.6 (br), 127.9, 124.2 (br). LRMS (EI) for $C_{13}H_9O_4N$ 243 (M^+ , 100), 152
20 (60); HRMS (EI) calcd 243.0531, obsd 243.0544. Anal. ($C_{13}H_9NO_4$) C, H, N.

C. 4-Nitro-3'-tert-butoxycarbonylbiphenyl (16).

 To a solution of 15 (1.215 g, 5 mmol) in 30 mL of methylene chloride was added oxalyl chloride
25 (0.65 mL, 7.45 mmol) and one drop of DMF. The mixture was stirred until no further bubbling was observed. The clear solution was evaporated to dryness to give the crude acid chloride. To
 another flask containing 7.0 mL of tert-butanol
30 was added n-BuLi (1.8 M in hexane, 2.8 mL, 5.04 mmol) under a water bath. The turbid solution was stirred for 5 min at room temperature and then the
 above acid chloride in 20 mL of THF was added through a dropping funnel. The mixture was
35 stirred overnight before the solvents were

evaporated. The residue was dissolved into methylene chloride and washed with 0.5 N NaOH. The organic layer was dried over MgSO_4 and evaporated. The residue was recrystallized from methanol to give pale yellow crystals (851 mg, 57%). m.p. 110.5-111.0°C. ^1H NMR (CDCl_3) δ 8.32 (d, 7.8 Hz, 2H), 8.24 (s, 1H), 8.06 (d, 7.7 Hz, 1H), 7.77 (m, 3H), 7.56 (t, 7.7 Hz, 1H), 1.63 (s, 9H). ^{13}C NMR (CDCl_3) δ 165.1, 147.1, 146.5, 138.7, 132.8, 131.0, 129.6, 129.0, 128.2, 127.8, 124.0, 81.4, 28.0, LRMS (EI) for $\text{C}_{17}\text{H}_{17}\text{O}_4\text{N}$ 299 (M^+ , 20), 243 (70), 266 (30), 152 (25; HRMS (EI) calcd 299.1157, obsd 299.1192. Anal. ($\text{C}_{17}\text{H}_{17}\text{NO}_4$) C, H, N.

D. N-Boc-S-trityl cysteinal (17).

To a solution of N-Boc-S-trityl cysteine (7.44 g, 16 mmol) in 85 mL of methylene chloride was added triethylamine (2.22 mL, 16 mmol) and N,O-dimethylhydroxylamine hydrochloride (1.57 g, 16.1 mmol). This mixture was cooled in an ice bath and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI, 3.08 g, 16.0 mmol) and HOBT (2.17 g, 16 mmol) was added. The mixture was stirred at 0°C for 1 hr and at room temperature for a further 10 hr. The mixture was extracted with methylene chloride and 0.5 N HCl. The organic layer was washed consecutively with 0.5 N HCl, concentrated NaHCO_3 and brine. The organic layer was dried and evaporated. The residue was purified by flash column chromatography (1.5:1=hexane:ethylacetate) to give a white foam (7.40 g, 91%). m.p. 59-60° (decomp). ^1H NMR (CDCl_3) δ 7.41 (m, 6H), 7.20-7.31 (m, 9H), 5.13 (d, 8.9 Hz, 1H), 4.76 (br s, 1H), 3.64 (s, 3H), 3.15 (s, 3H), 2.56 (dd, 4.7 and 12.1 Hz, 1H), 2.39 (dd, 7.8 and 12.1 Hz, 1H), 1.43 (s, 9H). ^{13}C NMR (CDCl_3) δ 170.7, 154.9, 144.2, 129.3, 127.6, 126.4,

79.3, 66.4, 61.2, 49.5, 33.8, 31.8, 28.1. This carboxyamide (2.02 g, 4.0 mmol) was dissolved in 30 mL of ether and cooled to -10°C. Lithium aluminum hydride (167 mg, 4.40 mmol) was added and the mixture was stirred for 15 min under the nitrogen. Then 40 mL of 0.5 N HCl was added and the solution was extracted with ether. The ether layer was washed with 0.5 N HCl and dried. The evaporation of solvents gave a white foam (1.80 g) which was used for further reaction without purification. The ¹H NMR spectrum of this compound was complex. The percentage of the aldehyde was about 65-70%, which was calculated according to the integration of the sharp singlet (δ 9.17) and the trityl peak (δ 7.40, m, 6H; 7.28, m, 9H). Lowering the temperature to -45°C did not improve the aldehyde percentage.

E. 4-N-[2(R)-tert-butoxycarbonylamino-3-triphenylmethylthiopropyl]amino-3'-tert-butoxycarbonylbiphenyl (18).

Compound 16 (768 mg, 2.56 mmol) was dissolved in THF. A catalytic amount of 10% Pd on activated carbon (78 mg) was added. The mixture was hydrogenated (40 psi) for 30 min. The black mixture was passed through a thin layer of celite and the pale yellow solution was evaporated. The residue was dissolved in 10 mL of methanol. To this solution was added 0.5 mL of acetic acid and a solution of the same equivalents of aldehyde 17 (according to the ¹H NMR determination) in 6 mL of methanol. Sodium cyanoborohydride (241 mg, 3.84 mmol, 1.5 eq) was added and the mixture was stirred overnight. After the evaporation of solvents, the residue was extracted with ethyl acetate and concentrated sodium bicarbonate. The organic layer was dried and evaporated. The

residue was purified by flash column chromatography (3.5:1-hexane:THF) to give a white foam (1.09 g, 61%). m.p. 75.0-76.0 °C(decomp).

[α]_D²⁵ = -2.13 (c=0.01, CH₃COOC₂H₅). ¹H NMR (CDCl₃) δ 8.14 (s, 1H), 7.86 (d, 7.7 Hz, 1H), 7.66 (d, 7.8 Hz, 1H), 7.40 (m, 9H), 7.22-7.30 (m, 9H), 6.61 (d, 8.5 Hz, 2H), 4.58 (d, 7.1 Hz, 1H), 3.83 (br m, 2H, Cys α proton and the amine), 3.12 (br m, 2H, CH₂N), 2.48 (br, m, 2H, CH₂S), 1.60 (s, 9H), 1.44 (s, 9H). ¹³C NMR (CDCl₃) δ 165.9, 155.6, 147.5, 144.4, 141.2, 132.3, 130.1, 129.5, 129.2, 128.5, 128.0, 127.9, 127.1, 126.8, 112.9, 80.9, 79.7, 67.0, 49.4, 47.1, 34.3, 28.3, 28.2 (expect 14 aromatic C, observed 13). Anal. (C₄₄H₄₈N₂O₄S·1.2H₂O) C, H, N, S.

F. 4-N-[2(R)-amino-3-mercaptopropyl]amino-3'-carboxybiphenyl (4).

Compound 18 (600 mg, 0.85 mmol) was dissolved in 2mL of TFA and 2 mL of methylene chloride. Triethylsilane was added dropwise to the deep brown mixture until the brown color had disappeared. The mixture was then kept at room temperature for 1 hr. Then solvents were evaporated and the residue was dried under vacuum. The solid was triturated with 30 mL of ether and 3mL of 3 N HCl in ether. The white precipitate was filtered and washed with ether to obtain a crude product (270 mg, 84%). This crude product was dissolved into 30 mL of dilute HCl solution (0.01N) and was lyophilized. Analytical HPLC showed the purity to be 95%. m.p. 105-106 °C (decomp). [α]_D²⁵ = +13.16 (c=0.01 in methanol. ¹J M<R (CD₃OD δ 8.18 (s, 1H), 7.89 (d, 7.7 Hz, 1H), 7.78 (d, 7.3 Hz, 1H), 7.49 (m, 3H), 6.82 (d, 8.5 Hz, 2H), 3.56 (m, 2H CHN and CH₂N), 3.42 (dd, 8.9 and 15.2 Hz, 1H, CH₂S). ¹³C NMR (D₂O and CD₃OD) δ

171.1, 147.1, 141.4, 131.9, 130.9, 130.1, 128.8, 128.5, 127.0, 115.2, 53.2, 45.4, 25.0 LRMS (FAB, glycerol) for $C_{16}H_{18}N_2O_2S$ (M+1) 303. Anal.

($C_{16}H_{18}N_2O_2S \cdot 2HCl$) C, H, N, S. Further purification
5 by preparative HPLC (Waters C-18, 40% acetonitrile, 60% water, 0.1% TFA, 40 min gradient) gave product 4 (120 mg) with a purity over 99.9%.

G. 3-Methyl-3'-tert-butoxycarbonylbiphenyl (19).

10 The coupling of 3-methylphenylboronic acid with 3-bromobenzoic acid methyl ester gave a 3-methyl-3'-methoxycarbonylbiphenyl (79% yield), which was then hydrolyzed to yield a 3-methyl-3'-carboxybiphenyl (97% yield). Compound 19 (an oil)
15 was prepared from this acid using the same method as for the preparation of compound 16 (65% yield). 1H NMR ($CDCl_3$) δ 8.21 (s, 1H), 7.95 (d, 7.8 Hz, 1H), 7.73 (d, 6.6 Hz, 1H), 7.46 (m, 3H), 7.35 (t, 7.5 Hz, 1H), 7.20 (d, 7.4 Hz, 1H), 2.43 (s, 3H), 1.62
20 (s, 9H). ^{13}C NMR ($CDCl_3$) δ 165.6, 141.3, 140.2, 138.3, 132.4, 140.0, 128.7, 128.5, 128.3, 128.0, 127.9, 124.2, 81.0, 28.1, 21.4. LRMS (EI) for $C_{18}H_{20}O_2$ 268 (M $^+$, 35), 212 (100), 195 (20); HRMS (EI) calcd 268.1463, obsd 268.1458.

25 H. 3-Azido-3'-tert-butoxycarbonylbiphenyl (20).

Compound 19 (2.18 g, 8.13 mmol) and N-bromosuccinimide (1.70 g, 9.50 mmol) was suspended
in 60 mL of CCl_4 . Dibenzoyl peroxide (20 mg) was added and the mixture was refluxed for 1.5 hr.
30 After removing the solid, the filtrate was washed with concentrated sodium bicarbonate and dried over sodium sulfate. 1H NMR showed the crude material contained 70% of monobrominated and 30% of dibrominated product. This material was
35 dissolved in 20 mL of DMSO and sodium azide (3.70

g, 57 mmol) was added. The mixture was heated to 80 °C for 4 hr before being poured into a mixture of methylene chloride and water. The organic layer was dried and evaporated. The residue was
5 purified by flash column chromatography (5% of ethyl acetate in hexane) to give 20 (2.14 g, 78%, two steps) as colorless oil. ¹H NMR (CDCl₃) δ 8.22 (s, 1H), 8.00 (d, 7.7 Hz, 1H), 7.76 (d, 8.2 Hz, 1H), 7.58 (m, 2H), 7.50 (m, 2H), 7.33 (d, 7.6 Hz,
10 1H), 4.43 (s, 2H), 1.62 (s, 9H). ¹³C NMR (CDCl₃) δ 165.2, 140.5, 140.3, 135.8, 132.3, 130.7, 129.1, 128.5, 128.2, 127.8, 127.1, 126.7, 126.6, 80.9, 54.3, 27.8.

15 I. N-Boc-S-trityl-cysteiny-3-aminomethyl-3'-tert-butoxycarbonylbiphenyl (21).

Compound 20 (0.75 g, 2.43 mmol) was dissolved in 30 mL of methanol. A catalytic amount of 5% palladium on barium sulfate (0.30 g) was added. The mixture was hydrogenated at 1 atm for 5 hr.
20 The catalyst was removed by filtration and the methanol was evaporated. This residue was dissolved in 40 mL of methylene chloride. N-Boc-S-trityl cysteine (1.12 g, 2.43 mmol) was added at 0°C followed by EDCI (1 eq) and HOBT (1 eq). The
25 mixture was stirred for 24 hr. After workup and evaporation of solvents, the residue was purified by flash column chromatography (hexane:ethyl acetate=3.2:1) to give 21 (570 mg, 44%). m.p. 84-86°C. ¹H NMR (CDCl₃) δ 8.17 (s, 1H), 7.95 (d, 7.7
30 Hz, 1H), 7.70 (d, 7.7 Hz, 1H), 7.50-7.30 (m, 9H), 7.30-7.10 (m, 11H), 6.44 (br, 1H), 4.86 (br, 1H), 4.45 (d, 4.0 Hz, 2H, CH₂Ph), 3.87 (br, 1H, Cys α H), 2.75 (dd, 7.2 and 12.8 Hz, 1H, CH₂S), 2.55 (dd, 5.3 and 12.8 Hz, 1H, CH₂S), 1.62 (s, 9H), 1.36 (s,
35 9H). Anal. (C₄₅H₄₈N₂O₅S) C, H, N, S.

J. Cysteinyl-3-aminomethyl-3'-carboxybiphenyl (11).

Compound 21 (150 mg) was deprotected using the same method as for the preparation of compound 4. Final purification by preparative HPLC gave 11 as a white solid (42 mg, 46%). m.p. 88-89 °C (decomp). ¹H MNR (CD₃OD) δ 8.26 (s, 1H), 8.01 (d, 7.7 Hz, 1H), 7.86 (d, 7.7 Hz, 1H), 7.64 (s, 1H), 7.56 (m, 2H), 7.46 (t, 7.6 Hz, 1H), 7.35 (d, 7.6 Hz, 1H), 4.53 (s, 2H), 4.00 (t, 5.2 Hz, 1H, Cys α H), 3.06 (dd, 14.6 and 5.2 Hz, 1H, CH₂S), 2.97 (dd, 14.6 and 6.8 Hz, 1H, CH₂S). LRMS (EI) for C₁₇H₁₈N₂O₃S 331 (M+1, 8), 281 (100), 226 (75), Anal. (C₁₇H₁₈N₂O₃S·HCl·0.6H₂O) C, H, N.

K. 3-N-[2(R)-tert-Butoxycarbonylamino-3-triphenylmethylthiopropyl]aminomethyl-3'-tert-butoxy-carboxybiphenyl (22).

The azide 20 (900 mg, 2.91 mmol) was dissolved in 20 mL of methanol. A catalytic amount of 5% Pd on barium sulfate (90 mg) was added. This mixture was hydrogenated at 1 atm overnight. The catalyst was removed and the methanol was evaporated. The remaining residue was dissolved in a mixture of 0.5 N HCl (20 mL) and ether (20 mL). The aqueous phase was neutralized with 1 N NaOH and extracted into methylene chloride. After the evaporation of solvents, a viscous oil was obtained (600 mg, 73%). ¹H NMR (CDCl₃) δ 8.22 (s, 1H), 7.97 (d, 7.8 Hz, 1H), 7.75 (d, 7.7 Hz, 1H), 7.57 (s, 1H), 7.50 (m, 2H), 7.43 (t, 7.7 Hz, 1H), 7.33 (d, 7.4 Hz, 1H), 3.96 (s, 2H), 1.62 (s, 9H), 1.46 (br s, 2H, NH₂). This amine (581 mg, 2.05 mmol) was dissolved in 10 mL of methanol and 0.5 mL of acetic acid before N-Boc-S-tritylcysteinyl (1eq, according to ¹H NMR determination of the aldehyde percentage)

was added. Sodium cyanoborohydride (193 mg, 1.50 eq) was added to the above solution and the mixture was stirred at room temperature overnight. After workup, the crude residue was purified by
5 flash column chromatography (1: 1 = ethyl acetate: hexane) to give a white foam (602 mg, 41%). m.p. 66-68 °C (decomp). ¹H NMR (CDCl₃) δ 8.21 (s, 1H), 7.96 (d, 7.7 Hz, 1H), 7.73 (d, 8.0 Hz, 1H), 7.37-7.51 (m, 10H), 7.15-7.31 (m, 10H), 4.69 (br
10 d, 1H), 3.75 (br s, 3H, PhCH₂N and Cys α H), 2.68 (dd, 6.0 and 12.3 Hz, 1H, CH₂S), 2.56 (dd, 5.5 and 12.3 Hz, 1H, CH₂S), 2.47 (m, 1H, CH₂N), 2.35 (m, 1H, CH₂N), 1.62 (s, 9H), 1.42 (s, 9H), 1.12 (br s, 1H, NH).

15 **L. 3-N-[2(R)-amino-3-mercaptopropyl] aminomethyl-3'-carboxybiphenyl (12).**

Compound 22 (480 mg, 0.672 mmol) was dissolved in a mixture of 2 mL of methylene chloride and 2 mL of trifluoroacetic acid.
20 Several drops of triethylsilane were added until the deep brown color had disappeared. This mixture was kept at room temperature for 1.5 hr, and then the solvents were evaporated, and the residue was dried under vacuum. The solid residue
25 was dissolved in 1 mL of acetic acid and 2 mL of HCl (1.7 M) in acetic acid. Finally 5 mL of HCl (3 M) in ether and 10 mL of ether were added. The white precipitate was washed with dry ether and dried to give a hydrochloride salt (215 mg, 81%).
30 ¹H NMR (D₂O) δ 8.16 (s, 1H), 7.94 (d, 7.7 Hz, 1H), 7.85 (d, 7.7 Hz, 1H), 7.70 (s, 2H), 7.55 (t, 7.8 Hz, 2H), 7.46 (d, 7.5 Hz, 1H), 4.36 (s, 2H, PhCH₂), 3.81 (m, 1H, Cys α H), 3.57 (dd, 5.7 and 13.7 Hz, 1H, CH₂N), 3.44 (dd, 6.5 and 13.7 Hz, 1H, CH₂N),
35 2.97 (dd, 5.3 and 15.1 Hz, 1H, CH₂S), 2.86 (dd, 5.9 and 15.1 Hz, 1H, CH₂S).

M. 2-Methoxy-4-nitro-3'-tert-butoxycarbonyl-biphenyl (23).

The coupling of 1-bromo-2-methoxy-4-nitrobenzene with 3-methylphenylboronic acid followed by the oxidation gave the 2-methoxy-4-nitro-3'-carboxybiphenyl. The reaction of acid chloride with lithium tert-butoxide gave 23 (3 steps, 35%). m.p. 88.0-88.5 °C. ¹H NMR (CDCl₃) δ 8.13 (s, 1H), 8.00 (d, 7.7 Hz, 1H), 7.89 (d, 8.3 Hz, 1H), 7.81 (s, 1H), 7.69 (d, 7.7 Hz, 1H), 7.48 (m, 2H), 3.90 (s, 3H), 1.60 (s, 9H). ¹³C NMR (CDCl₃) δ 165.2, 156.7, 148.0, 136.3, 136.2, 133.2, 132.0, 130.8, 130.1, 129.0, 127.9, 115.8, 106.0, 81.1, 55.9, 27.9. LRMS (EI) for C₁₈H₁₉NO₅ 329 (M⁺, 30), 273 (100).

N. 2-Methoxy-4-N-[2(R)-N-tert-butoxycarbonylamino-3-triphenylmethylthiopropyl]amino-3'-tert-butoxycarbonylbiphenyl (24).

Compound 24 was prepared using the same method as for the preparation of compound 18 (yield 63%). m.p. 76.0-77.0 °C (decomp). [α]_D²⁵ = -11.25 (c=0.01, CH₃COOC₂H₅). ¹H NMR (CDCl₃) δ 8.09 (s, 1H), 7.86 (d, 7.0 Hz, 1H), 7.65 (d, 7.0 Hz, 1H), 7.37 (t, 7.7 Hz, 1H), 7.43 (m, 6H), 7.21-7.32 (m, 9H), 7.11 (d, 8.1 Hz, 1H), 6.21 (s, 1H), 6.18 (d, 8.1 Hz, 1H), 4.58 (d, 6.1 Hz, 1H), 3.86 (br s, 1H), 3.76 (s and m, 4H), 3.14 (br d, 4.9 Hz, 2H), 2.49 (br d, 5.1 Hz, 2H), 1.59 (s, 9H), 1.43 (s, 9H). ¹³C NMR (CDCl₃) δ 165.9, 157.3, 155.5, 148.8, 144.3, 138.9, 133.3, 131.5, 131.2, 130.0, 129.4, 127.8, 127.5, 126.7, 118.7, 104.7, 96.2, 80.5, 79.4, 66.8, 55.2, 49.3, 47.0, 34.1, 28.2, 28.1. Anal. (C₄₅H₅₀N₂O₅S) C, H, N, S.

O. 2-Methoxy-4-N-[2(R)-amino-3-mercaptopropyl]amino-3'-carboxybiphenyl (10).

Compound 10 was obtained from the deprotection of compound 24. m.p. 120-121 °C (decomp). $[\alpha]^{25}_D = +12.62$ ($c = 0.01$, in methanol). ^1H NMR (CD_3OD) δ 8.09 (s, 1H), 7.89 (d, 7.8 Hz, 1H), 7.67 (d, 7.8 Hz, 1H), 7.43 (t, 7.7 Hz, 1H), 7.20 (d, 8.1 Hz, 1H), 6.56 (s, 1H), 6.53 (d, 8.1 Hz, 1H), 3.81 (s, 3H), 3.60 (m, 2H, Cys α H and CH_2N), 3.48 (m, 1H, CH_2N), 2.96 (dd, 4.9 and 13.7 Hz, 1H, CH_2S), 2.86 (dd, 5.4 and 13.7 Hz, 1H, CH_2S). ^{13}C NMR (D_2O and CD_3OD) δ 171.1, 158.2, 149.3, 139.7, 135.1, 132.2, 131.1, 130.4, 129.4, 128.4, 120.5, 106.2 (broad, due to deuterium exchange), 98.8, 56.3, 53.4, 45.1, 24.9. LRMS (EI) for $\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_3\text{S}$ 332 (M^+). Anal. ($\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}_3\text{S} \cdot 1.2\text{HCl} \cdot \text{H}_2\text{O}$) C, H, N, S.

15 **P. Cysteiny-4-amino-3'-carboxybiphenyl (6).**

Compound 6 was purified through preparative HPLC. Purity was shown to be over 99%. m.p. 120.0-121.0 °C. ^1H NMR (CD_3OD) δ 8.25 (s, 1H), 7.98 (d, 7.6 Hz, 1H), 7.84 (d, 7.7 Hz, 1H), 7.74 (d, 7.0 Hz, 2H), 7.54 (t, 7.7 Hz, 1H), 7.66 (d, 8.6 Hz, 2H), 4.16 (q, 5.0 Hz, 1H), 3.19 (dd, 5.20 and 14.8 Hz, 1H), 3.07 (dd, 7.7 and 14.7 Hz, 1H); ^{13}C NMR (CD_3OD) δ 169.8, 166.7, 141.9, 138.7, 137.8, 132.6, 132.2, 130.2, 129.5, 128.8, 128.5, 121.6, 56.9, 26.4. LRMS (EI) for $\text{C}_{16}\text{H}_{16}\text{O}_3\text{N}_2\text{S}$ 316 (M^+ , 25), 213 (100); HRMS (EI) calcd 316.0882, obsd 316.0867. Anal. ($\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_3\text{S} \cdot \text{CF}_3\text{COOH} \cdot \text{H}_2\text{O}$) C, H, N.

30 **Q. 4-N-[2(R)-Amino-3-mercaptopropyl]amino-2'-carboxybiphenyl (3).**

m.p. 129-130 °C (decomp). $[\alpha]^{25}_D = +12.58$ ($c = 0.01$, CH_3OH). ^1H NMR (CD_3OD) δ 7.73 (d, 7.6 Hz, 1H), 7.50 (d, 7.6 Hz, 1H), 7.35 (m, 2H), 7.21 (d, 8.5 Hz, 2H), 6.86 (d, 8.5 Hz, 2H), 3.58 (m, 2H), 3.46 (m, 1H), 2.97 (dd, 4.8 and 14.6 Hz, 1H), 2.86 (dd, 5.4 and 14.6 Hz, 1H). ^{13}C NMR (D_2O and CD_3OD) δ

174.3, 146.3, 141.9, 132.8, 132.7, 131.4, 130.6, 130.2, 127.9, 115.3, 52.9, 45.8, 25.0. LRMS (FAB, glycerol) for $C_{16}H_{18}N_2O_2S$ (M+1) 303. Anal. ($C_{16}H_{18}N_2O_2S \cdot 1.6HCl$) C, H, N, S.

5 R. 4-N-[2(R)-Amino-3-mercaptopropyl]amino-4'-carboxybiphenyl (5).

m.p. 260 °C (decomp). $[\alpha]^{25}_D = +12.20$ (c=0.01, CH_3OH). 1H NMR (CD_3OD) δ 8.03 (d, 8.5 Hz, 2H), 7.66 (d, 8.4 Hz, 2H), 7.56 (d, 8.4 Hz, 2H), 6.85 (d, 8.5 Hz, 2H), 3.57 (m, 2H), 3.45 (m, 1H), 2.98 (dd, 4.8 and 14.5 Hz, 1H), 2.85 (dd, 5.7 and 14.5 Hz, 1H). ^{13}C NMR (D_2O and CD_3OO) δ 169.8, 146.4, 146.1, 133.6, 131.4, 129.8, 129.4, 127.1, 117.0, 53.3, 47.2, 25.5. LRMS (EI) for $C_{16}H_{18}O_2N_2S$ 302 (M⁺, 15), 285 (15), 226 (100), 213 (50). HRMS (EI) calcd 302.1088, obsd 302.1089. Anal. ($C_{16}H_{18}N_2O_2S \cdot 2HCl$) C, H, N.

S. 4-N-[2(R)Amino-3-Mercaptopropyl]aminobiphenyl (7).

20 m.p. 216 °C (decomp). $[\alpha]^{25}_D = +13.27$ (c=0.01, CH_3OH). 1H NMR (CD_3OD) α 7.54 (m, 4H), 7.39 (m, 2H), 7.26 (m, 1H), 6.82 (br s, 2H), 3.56 (br m, 2H), 3.45 (m, 1H), 2.98 (m, 1H), 2.87 (m, 1H); ^{13}C NMR (CD_3OD) δ 144.8, 141.8, 135.7, 129.9, 129.1, 127.8, 127.4, 117.4, 53.2, 47.6, 25.5. LRMS (EI) for $C_{15}H_{16}N_2S$ 258 (M⁺, 15), 182 (100); HRMS (EI) calcd 258.1190, obsd 258.1183. Anal. ($C_{15}H_{16}N_2S \cdot 1.6HCl$) C, H, N, S.

30 T. 4-N-[2(R)-Amino-3-mercaptopropyl]amino-3'-methylbiphenyl (8).

1H NMR (CD_3OD) δ 7.50 (d, 8.2 Hz, 2H), 7.35 (m, 2H), 7.27 (t, 7.6 Hz, 1H), 7.08 (d, 7.3 Hz, 1H), 6.95 (d, 8.2 Hz, 2H), 3.60 (m, 2H), 3.46 (m, 1H), 2.99 (dd, 4.9 and 14.6 Hz, 1H), 2.88 (dd, 5.5

and 14.6 Hz, 1H), 2.37 (s, 3H). LRMS (EI) for $C_{16}H_{20}N_2S$ 272 (M^+ , 15), 196 (100). HRMS (EI) calcd 272.1341, obsd 272.1347.

5 U. 4-N-[2(R)-Amino-3-mercaptopropyl]amino-3'-methoxycarbonylbiphenyl (9).

m.p. 86-89 °C (decomp). 1H NMR (CD_3OD) δ 8.18 (s, 1H), 7.90 (d, 7.7 Hz, 1H), 7.79 (d, 6.6 Hz, 1H), 7.55 (d, 8.6 Hz, 2H), 7.49 (t, 7.7 Hz, 1H), 7.01 (d, 8.6 Hz, 2H), 3.92 (s, 3H), 3.543.65 (m, 2H), 3.44-3.52 (m, 1H), 2.97 (dd, 4.7 and 14.7 Hz, 1H), 2.88 (dd, 5.5 and 14.7 Hz, 1H). LRMS (EI) for $C_{17}H_{20}O_2N_2S$ 316 (M^+ , 15), 299 (20), 240 (100). HRMS (EI) calcd 316.1240, obsd 316.1239. Anal. ($C_{17}H_{20}N_2O_2S \cdot 2HCl$) C, H, N, S.

Table 2

Table of Microanalysis Data

Compd	Formulae	C% (cal, obs)	H% (cal, obs)	N% (cal, obs)	S% (cal, obs)
3	$C_{16}H_{18}N_2O_2S \cdot 1.6HCl$	53.27 (53.41)	5.44 (5.78)	7.77 (7.35)	8.87 (8.47)
4	$C_{16}H_{18}N_2O_2S \cdot 2HCl$	51.20 (51.60)	5.37 (5.30)	7.47 (7.07)	8.53 (8.22)
5	$C_{16}H_{18}N_2O_2S \cdot 2HCl$	51.20 (51.62)	5.37 (5.56)	7.47 (7.00)	
6	$C_{16}H_{16}N_2O_3S \cdot CF_3COOH \cdot H_2O$	48.21 (48.24)	4.24 (4.20)	6.25 (6.32)	
7	$C_{15}H_{18}N_2S \cdot 1.6HCl$	56.89 (57.04)	6.19 (6.46)	8.85 (8.74)	10.11 (10.03)
9	$C_{17}H_{20}N_2O_2S \cdot 2HCl$	52.44 (52.84)	5.65 (5.92)	7.19 (7.37)	8.22 (8.53)
10	$C_{17}H_{20}N_2O_3S \cdot 1.2HCl \cdot H_2O$	51.80 (51.91)	5.89 (5.96)	7.11 (6.81)	8.12 (7.77)
11	$C_{17}H_{18}N_2O_3S \cdot HCl \cdot 0.6H_2O$	54.07 (54.11)	5.35 (5.39)	7.42 (7.35)	
15	$C_{13}H_9NO_4$	64.19 (64.05)	3.70 (3.75)	5.76 (5.80)	
16	$C_{17}H_{17}NO_4$	68.23 (68.07)	5.68 (5.73)	4.68 (4.64)	
18	$C_{44}H_{48}N_2O_4S \cdot 1.2H_2O$	73.17 (72.82)	6.98 (6.83)	3.88 (3.87)	4.43 (4.50)
21	$C_{45}H_{48}N_2O_5S$	74.14 (73.74)	6.64 (6.74)	3.84 (3.79)	4.39 (4.32)
24	$C_{45}H_{50}N_2O_5S$	73.97 (73.72)	6.85 (7.04)	3.83 (3.66)	4.38 (4.32)

Table 3
Examples of Peptidomimetics of the Invention

	<u>Compound</u>	<u>Structure</u>
2	FTI-232	Cys-4-aminobenzoyl-Met
5	2a FTI-249	red.Cys-4-aminobenzoyl-Met
	3 FTI-273	red.Cys-4-amino-2'-carboxybiphenyl
	4 FTI-265	red.Cys-4-amino-3'-carboxybiphenyl
	5 FTI-271	red.Cys-4-amino-4'-carboxybiphenyl
	6 FTI-278	Cys-4-amino-3'-carboxybiphenyl
10	7 FTI-268	red.Cys-4-aminobiphenyl
	8 FTI-263	red.Cys-4-amino-3'-methylbiphenyl
	9 FTI-259	red.Cys-4-amino-3'-carboxymethylbiphenyl
	10 FTI-281	red.Cys-4-amino-2-OMe-3'-carboxybiphenyl
	11 FTI-285	red.Cys-4-amino-2-phenyl-3'-methylbiphenyl
15	12 FTI-238	Cys-3-aminomethyl-3'-carboxybiphenyl
	FTI-283	red.Cys-3-aminomethyl-3'-carboxybiphenyl
	FTI-282	(DL)4-(2,3-diaminopropyl)-amino-3' carboxybiphenyl
	FTI-288	red.Cys-4-amino-2-OPr-3'-carboxybiphenyl
	FTI-289	red.Cys-4-amino-2-phenyl-3'-carboxybiphenyl
20	FTI-291	4-(3-aminoalanyl)-amino-3'-carboxybiphenyl
	FTI-292	(L)4(2,3-diaminopropyl)-amino-3'-carboxybiphenyl
	FTI-295	4-(Ethylsulfonyl-3-aminoalanyl)-amino-3'-carboxy- biphenyl
25	FTI-296	4-(Vinylsulfonyl-3-aminoalanyl)-amino-3'- carboxybiphenyl
	FTI-2102	red.Cys-4-amino-3'-tetrazolylbiphenyl

Number designations used for compounds of the invention discussed below is shown in Table 3.

EXAMPLE 22

FTase and GGTase I Activity Assay

5 Human Burkitt lymphoma (Daudi) cells (ATCC, Rockville, MD) were grown in suspension in RPMI 1640 medium containing 10% fetal bovine serum (FBS) and 1% Pen-Strep in a humidified 10% CO₂ incubator at 37°C. The cells were harvested and
10 sonicated in 50 mM Tris, pH 7.5, 1 mM EDTA, 1 mM EGTA, 25 µg/ml leupeptin, 1 mM phenylmethylsulfonyl fluoride. Homogenates were then spun at 12,000 x g and the resulting supernatant further spun at 60,000 x g. The
15 supernatant was assayed for both FTase and GGTase I. Briefly, 100 µg of the supernatants was incubated in 50 mM Tris, pH 7.5, 50 µM ZnCl₂, 20 mM KCl, 3 mM MgCl₂, and 1 mM DTT. For FTase assays, the reaction was incubated at 37°C for 30 minutes
20 with recombinant H-Ras-CVLS (11 µM) and [³H] FPP (625 nM; 16.3 Ci/mmol). For GGTase assays, the reaction was also incubated for 30 minutes at 37°C but with recombinant H-Ras-CVLL (5 µM) and [³H] GGPP (525 nM; 19.0 Ci/mmol). The reaction was
25 stopped and passed through glass fiber filters to separate free and incorporated label. For inhibition studies, the peptidomimetics were premixed with FTase or GGTase I prior to adding the remainder of the reaction mixture.
30 Recombinant H-Ras-CVLS was prepared as described previously (26) from bacteria (31). Recombinant H-Ras-CVLL was prepared from bacteria (32).

EXAMPLE 23

Peptidomimetics Farnesylation Assay

The ability of human Burkitt lymphoma (Daudi) FTase to farnesylate peptides and peptidomimetics was determined as described previously (34, 35). Briefly, 25 μ l of reaction mixture containing 50 μ g of 60,000 xg supernatants and 20 μ M peptidomimetic in 50 mM Tris, pH 7.5, 50 μ M ZnCl₂, 20 mM KCl, 3 mM MgCl₂, 1 mM DTT and 0.2% octyl β -D-glucoside was incubated for 30 minutes at 37°C, then spotted onto silica gel G TLC sheets (20 x 20 cm, Brinkmann Instruments), and developed with n-propanol/5 N ammonium hydroxide/water (6:1:1). The dried sheets were sprayed with En³Hance (DuPont NEN) and exposed to x-ray film for detection of [³H] farnesylated products.

EXAMPLE 24

Ras and Rap1A Processing Assay

EJ3 cells were treated with peptidomimetics or vehicle for 20-24 h. Cells were lysed in lysis buffer (10 mM Na₂HPO₄, pH 7.25, 150 mM NaCl, 0.1% sodium dodecyl sulfate, 1% Triton X-100, 12 mM sodium deoxycholate, 1 mM NaF, 0.2% NaN₃, 2 mM PMSF, 25 μ g/ml leupeptin) and the lysates were cleared by spinning at 13,000 rpm for 15 minutes. Ras protein was immunoprecipitated overnight at 4°C with 50 μ g of anti-Ras antibody (Y13-259; hybridoma from ATCC, Rockville, MD) along with 30 μ l Protein A-agarose goat anti-rat IgG complex (Oncogene Science, Uniondale, NY). Immunoprecipitates were washed 4 times with lysis buffer and the bound proteins were released by heating for 5 minutes in 40 μ l SDS-PAGE sample buffer and subsequently electrophoresed on a 12.5% SDS-PAGE. Proteins were transferred onto nitrocellulose and subsequently blocked with 5%

non-fat dry milk in PBS (containing 1% Tween 20 (PBS-T) and probed with Y13-259 (50 µg/ml in 3% non-fat dry milk in PBS-T). Positive antibody reactions were visualized using peroxidase-conjugated goat anti-rat IgG (Oncogene Science, Uniondale, NY) and an enhanced chemiluminescence detection system (ECL; Amersham).

For Rap1A processing assays, 50 µg of cell lysates were electrophoresed as described above for Ras processing and transferred to nitrocellulose. These membranes were then blocked with 5% milk in Tris-buffered saline, pH 8.0, containing 0.5% Tween-20 and probed with anti-Rap1A (1 µg/ml in 5% milk/TBS-T; Santa Cruz Biotechnology, Santa Cruz, CA). Antibody reactions were visualized using peroxidase-conjugated goat anti-rabbit IgG (Oncogene) and ECL chemiluminescence as described above.

Structural Modeling (Figure 13)

The calculation of the energy minimized conformations was carried out using the AMBER force field within the MacroModel program, version 3.5a.

EXAMPLE 25

The potency of the peptidomimetics of Figure 12 and Table 3 for inhibiting partially purified FTase was evaluated by determining their ability to inhibit the transfer of farnesyl to recombinant H-Ras as described above. The results are summarized in Table 4, which indicates the IC₅₀s obtained for FTase activity and GGTase-I activity, and the selectivity for a number of peptidomimetics of the invention. The IC₅₀ values given in Table 4 represent inhibition of FTase and GGTase I in vitro by the listed compounds.

Table 4

In vitro Activity of CAAX Mimetic Inhibitors of FTase

Inhibitor	FTase IC ₅₀ (nM)	GGTase-I IC ₅₀ (nM)	Selectivity	Substrate
FTI-232	150	1500	10	
FTI-249	300	4400	15	
FTI-273	543 (3) ^a	140,000 (2) ^a	258	nd ^b
FTI-265	114 (10)	100,000 (6)	877	no
FTI-271	4575 (4)	>100,000 (2)	>22	no
FTI-278	13,500 (2)	100,000 (2)	7	nd
FTI-268	1,070 (3)	>100,000 (3)	93	no
FTI-263	710 (3)	>100,000 (3)	141	no
FTI-259	917 (3)	>100,000 (3)	109	no
FTI-281	40 (6)	43,600 (5)	1090	nd
FTI-238	100,000 (2)	>100,000 (3)	>1	no
FTI-283	11,000 (1)	35,000 (2)	3	nd
FTI-285	2075 (4)	8500 (2)	4	
FTI-282	50,000 (1)	>>100,000 (1)	>2	
FTI-288	41 (6)	2375 (4)	59	
FTI-289	16 (5)	643 (4)	40	
FTI-291	210,000 (1)		0	
FTI-292	60,000 (1)		0	
FTI-295	200,000 (1)	>1,000,000 (1)	>5	
FTI-296	430,000 (1)	>>1,000,000 (1)	>2	
FTI-2102	30 (1)	5,000 (1)	187	

^aNumbers in parentheses indicate number of determinations. Where no number is given, at least two determinations were made.

^bnd indicates not determined.

The results obtained showed that compound 2, i.e. Cys-4ABA-Met (1-10 μ M) inhibited FTase in a concentration-dependent manner with an IC_{50} of 150 nM (FTI-232, Table 4). This value is similar to the previously reported IC_{50} values for CVIM and Cys-4ABA-Met (35). Reduction of the amide bond between cysteine and aminobenzoic acid gave the red-Cys-4ABA-Met (2a, FTI-249) which had an IC_{50} of 300 nM. However, replacing the methionine and the C-terminal amide bond in (2a) by another aromatic ring to obtain the biphenyl-based peptidomimetic (4) improved potency by twofold (FTI-265, Table 4). Peptidomimetic 4 had an IC_{50} of 114 nM towards partially purified FTase from human Burkitt lymphoma cells and 50 nM towards rat brain FTase purified to homogeneity. Thus, despite major structural differences between the compound CVIM (1) and 4, the latter (4) retained the potent FTase inhibitory activity of the tetrapeptide CVIM (1) and the peptide mimetics 2 and 2a.

As noted, Figures 14A and 14B graphically illustrate the results of FTase and GGTase I inhibition studies. In these studies, partially purified FTase and GGTase I were incubated with the peptidomimetics to be tested and their ability to transfer [3 H] farnesyl to H-Ras-CVLS (FTase) and [3 H] geranylgeranyl to H-Ras CVLL (CCTase I) was determined as described. Figure 14A shows FTase inhibition by: \square , (4) and \blacksquare , (5) while Figure 14B plots FTase (\square) and GGTase I (\blacksquare) inhibition by (4). Each curve is representative of at least four independent experiments.

Geranylgeranylation is a more common protein prenylation than farnesylation (49). It is, therefore, advantageous for CAAX peptidomimetics targeting farnesylation to have high selectivity towards inhibiting FTase compared to GGTase. In

the CAAX tetrapeptides, the X position determines whether the cysteine thiol will be farnesylated by FTase or geranylgeranylated by GGTase I. Those proteins or peptides with Leu or Ile at the X position are geranylgeranylated. As shown in Table 4, the present compounds do not significantly inhibit GGTase I and demonstrate much greater selectivity for FTase.

Figure 14B shows that compound 4, which is a potent FTase inhibitor, is a very poor GGTase I inhibitor. The ability of compound 4 to inhibit the transfer of geranylgeranyl to Ras-CVLL ($IC_{50} = 100,000$ nM) was found to be 877-fold less than that of 4 to inhibit the transfer of farnesyl to Ras-CVLS ($IC_{50} = 114$ nM) (Table 4). This selectivity was much more pronounced than in the peptidomimetics 2 and 2a which were more selective for FTase relative to GGTase I by only 10 and 15-fold, respectively. It is also noted that the free carboxylate of compound 4 is not responsible for this selectivity since replacement of this group by a methyl in compound 8 did not increase affinity towards GGTase I (Table 4). These results indicate that the FTase and GGTase I binding sites are quite different and that differences between Leu, Ile and Met side chains cannot be the only predictors of selectivity. Regardless of the explanation, it is clear that the compounds of the invention are much more selective to inhibition of FTase.

Besides having poor cellular uptake and being rapidly degraded, another disadvantage of natural CAAX peptides is that they are farnesylated by FTase. This results in metabolic inactivation since farnesylated CAAX derivatives are no longer inhibitors of FTase (34). Figure 15 shows that the natural peptide CVLS (carboxyl terminal CAAX

of H-Ras) is farnesylated by FTase from Burkitt lymphoma cells. Replacing the tripeptide VLS with 4-amino-3'-hydroxycarbonylbiphenyl, as in 4 did not affect potency towards FTase inhibition (Table 4) but prevented farnesylation of the cysteine thiol (Figure 15). None of the peptidomimetics of the invention is metabolically-inactivated by FTase (Figure 15). Thus, although AAX tripeptides are not necessary for potent FTase inhibition, they appear to be required for farnesylation.

With reference to Figure 15, it is to be noted that the transfer of [³H] farnesyl to peptides and peptidomimetic by FTase was determined by silica G TLC as described below. FPP, F-peptide, and ORIGIN designate farnesyl pyrophosphate, farnesylated peptide and origin, respectively. Figure 15 shows: Lane 1, FPP only; lane 2, FPP and CVLS but no FTase; lane 3, FPP and FTase but not peptide. Lanes 4-9 all contained FTase and FPP with lane 4, CVIM; lane 5, CVLS; lane 6, compound 2a; lane 7, compound 4; lane 8, compound 5; lane 9, compound 8. The results shown indicate that the compounds of the invention are not farnesylated in contrast to the CAAX compounds. Data given are representative of two independent experiments.

The foregoing results show that the novel peptidomimetics described herein have two very important features, namely, they are potent FTase inhibitors, and they are resistant to metabolic inactivation by FTase. Another important feature is that the present compounds inhibit Ras processing in whole cells. This is shown by the following with reference to Figure 16 which illustrates Ras and Rap1A processing. To this end, Ras transformed 3T3 cells were treated with inhibitors, lysed and the lysate A)

immunoprecipitated with anti-Ras antibody or B) separated by SDS-PAGE. Immunoprecipitates from A) were separated by SDS-PAGE and blotted with anti-Ras antibody whereas samples from B) were blotted with anti-Rap1A antibody as described hereafter. Figure 16 shows: Lane 1, control; lane 2, lovastatin; lane 3, reduced 2a (200 μ M); lane 4, 4 (100 μ M); lane 5, 4 (50 μ M); lane 6, 4 (25 μ M); lane 7, 5; lane 8, 8. Data are representative of 3 independent experiments. Farnesylated Ras runs faster than unprocessed Ras on SDS-PAGE (23-25, 28, 29). Figure 16A (lane 1) shows that cells treated with vehicle contain only processed Ras whereas cells treated with lovastatin (lane 2) contained both processed and unprocessed Ras indicating that lovastatin inhibited Ras processing. Lovastatin, an HMG-CoA reductase inhibitor which inhibits the biosynthesis of farnesylpyrophosphate and geranylgeranylpyrophosphate, is used routinely as a positive control for inhibition of processing of both geranylgeranylated and farnesylated proteins (36, 37, 39, 40, 51). Cells treated with reduced Cys-4ABA-Met 3 in its free carboxylate forms did not inhibit Ras processing. However, in contrast, the corresponding methyl ester of 2a (200 μ M) inhibited FTase (Figure 16A, lane 3). This is consistent with previous work that showed that neutralization of the carboxylate of CAAX peptides enhances their ability to inhibit Ras processing (37, 40, 51). Although compound 4 has a free carboxylate negative charge, it was able to enter cells and potently inhibit Ras processing (lane 4, 100 μ M compound 4). It was found that compound 4 inhibited Ras processing with concentrations as low as 50 μ M (lane 5), whereas its corresponding parent compound 2a did not inhibit Ras processing

at concentrations as high as 200 μ M. Compound 4 was as potent as the methylester of its parent compound (2a) (Figure 16A, lane 3). Furthermore, 4 appears to be the first CAAX peptidomimetic that effectively inhibits Ras processing in whole cells directly without relying on cellular enzymes for activation. The hydrophobic character of the biphenyl group apparently compensates for the free carboxylate negative charge thus allowing the peptidomimetic to penetrate membranes and promoting its cellular uptake.

The selectivity of the present Ras farnesylation inhibitors has also been investigated by determining their ability to inhibit processing of Rap1A, a small G-protein that is geranylgeranylated (49, 50). Cells were treated with lovastatin or peptidomimetics exactly as described for Ras processing experiments. Lysates were then separated by SDS-PAGE and immunoblotted with anti-Rap1A antibody as described below. Control cells contained only the geranylgeranylated Rap1A (Figure 16B, lane 1) whereas lovastatin-treated cells contained both processed and unprocessed forms of Rap1A indicating, as expected, that lovastatin inhibited the processing of Rap1A (Figure 16B, lane 2). Compound 4, which inhibited Ras processing, was not able to inhibit Rap1A geranylgeranylation (Figure 16B, lanes 4-6). Compounds 5 and 8 also did not inhibit Rap1A processing (Figure 16B, lanes 7 and 8).

Structures for a number of compounds of the invention are summarized in Table 3, and their relative effectiveness in inhibiting FTTase and GGTase shown in Table 4. The activity of the inhibitors is reported in Table 4 as IC_{50} values, the concentration at which FTase or GGTase I

activity was inhibited by 50%. Some of the inhibitors were further characterized for their ability to serve as substrates for farneylsation by thin layer chromatography, as shown in Table 4 (41).

The published sequence dependence studies on FTase have shown a strong preference for methionine in the terminal position of CAAX. In the compounds of the present invention, no methionine residue is present and the tripeptide AAX is completely replaced by a simple hydrophobic moiety. The most potent inhibitor in the CAAX series is Cys-Ile-Phe-Met (18, 22) with an IC_{50} value of 30 nM. Peptidomimetic inhibitor 10 is as potent as CIFM despite the large difference between their structures. These results confirm the hydrophobic strategy for AAX replacement according to the invention.

As previously reported, the incorporation of an aromatic amino acid into the A_2 position of CA_1A_2X (such as CIFM) prevents the tetrapeptide from serving as a substrate for farneylsation (22). Table 4 shows that the designed non-peptide CAAX mimetics (such as compound 4) are not substrates for farnesylation. This lack of farnesylation by FTase may be due to the inhibitor binding to the enzyme in a conformation that does not permit farnesyl transfer to the thiol group.

III. Geranylgeranyl tranferase Inhibitors

The carboxyl terminal CAAX tetrapeptide of Ras is a substrate for FTase and serves as a target for designing inhibitors of this enzyme with potential anticancer activity (33). Our earlier application describes a highly potent ($IC_{50}=500$ pM) inhibitor of FTase, FTI-276 (Fig. 17)

(66). Its cell-permeable methyl ester FTI-277 inhibits H-Ras processing in whole cells with an IC_{50} of 100 nM (66). Furthermore, FTI-276 is highly selective (100-fold) for FTase over GGTase I (Table 5).

TABLE 5

	In Vitro (IC_{50} , nM)		In Vivo Processing (IC_{50} , μ M)			
	FTase	GGTase I	H-Ras	K-Ras	Rap1A	
FTI-276	0.5	50	FTI-277	0.1	10	50
GGTI-287	25	5	GGTI-286	>30	2	2
GGTI-297	270	40	GGTI-298	>20	3	3

EXAMPLE 26

Synthesis of FTase and GGTase I Inhibitors

Peptidomimetics FTI-276 and FTI-277 were prepared as described above. GGTase I inhibitors GGTI-287 and GGTI-286 were prepared from 2-phenyl-4-nitrobenzoic acid (66) by reaction with L-leucine methyl ester followed by reduction with stannous chloride. The resulting 4-amino-2-phenylbenzoyl leucine methyl ester was reacted with N-Boc-S-trityl-cysteinal and deprotected by procedures similar to those described for the FTase inhibitors (66) to give GGTI-286 and GGTI-287 as their hydrochloride salts.

A. 4-Amino-2-phenylbenzoyl-(S)-methionine methyl ester hydrochloride

To a mixture of 70 mL of acetone and 85 mL of water was added 2-bromo-4-nitrotoluene 6.84 g (30 mmol), phenylboronic acid 3.84 g (31.5 mmol), potassium carbonate 10.35 g (75 mmol) and palladium acetate 336 mg (1.5 mmol). The mixture was refluxed for 10 hr and then extracted with ether and dilute hydrochloric acid. After evaporating solvents, the solid residue was recrystallized from methanol to give 5.64 g of 4-

nitro-2-phenyltoluene (88% yield). ^1H NMR (CDCl_3) δ 8.09-8.11 (m, 2H), 7.40-7.49 (m, 4H), 7.30-7.33 (m, 2H), 2.37 (s, 3H).

5 The above 4-nitro-2-phenyltoluene (4.46 g, 21 mmol) was suspended in 21 mL of pyridine and 42 mL of water. The mixture was heated to boiling followed by addition of potassium permanganate (19.8, 126 mmol). The mixture was refluxed for 2 hr and then filtered to remove the solids. The
10 filtrate was acidified with 6N HCl to give 4.48 g of 4-nitro-2-phenylbenzoic acid (89% yield). ^1H NMR (CDCl_3) δ 8.25-8.33 (m, 2H), 8.08 (d, 8.9 Hz, 1H), 7.41-7.51 (m, 3H), 7.31-7.39 (m, 2H).

15 The above 4-nitro-2-phenylbenzoic acid (2.43 g, 10 mmol) was suspended in 50 mL of methylene chloride. To this solution was added (L)-methionine methyl ester hydrochloride (2.0 g, 10 mmol), triethylamine (1.38 mL, 10 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
20 hydrochloride (EDCI, 2.01 g, 10.5 mmol), 1-hydroxybenzotriazole (HOBT, 1.35 g, 10 mmol). The mixture was stirred for 12 hr and then extracted with methylene chloride and 1N hydrochloric acid. After the evaporation of solvents, the residue was
25 recrystallized from ethyl acetate and hexane to give 3.22 g of 4-nitro-2-phenylbenzoyl-(L)-methionine methyl ester (yield 83%). ^1H NMR (CDCl_3) δ 8.24-8.28 (m, 2H), 7.85 (d, 8.9 Hz, 1H), 7.43-7.52 (m, 5H), 6.01 (d, 7.5 Hz, 1H), 4.69
30 (ddd, 1H), 3.68 (s, 3H), 2.05 (m, 2H), 1.98 (s, 3H), 1.88-1.96 (m, 1H), 1.72-1.81 (m, 1H).

The 4-nitro-2-phenylbenzoyl-(L)-methionine methyl ester (3.04 g, 7.83 mmol) was dissolved
35 into 100 mL of ethyl acetate followed by the addition of stannous chloride hydrate (8.84 g, 39 mmol). The mixture was refluxed for 2 hr and then extracted with a mixture of ethyl acetate and

concentrated sodium bicarbonate. After the evaporation of solvents, the residue was dissolved in methylene chloride followed by addition of 3N hydrogen chloride in ether. The solid was
5 filtered and dried to give 2.96 g of 4-amino-2-phenylbenzoyl-(L)-methionine methyl ester hydrochloride (yield 96%). ¹H NMR (CD₃OD) δ 7.65 (d, 8.1 Hz, 1H), 7.39-7.46 (m, 7H), 4.53 (dd, 4.3 and 9.5 Hz, 1H), 3.69 (s, 3H), 2.15-2.23 (m, 1H),
10 2.00 (s, 3H), 1.93-2.11 (m, 2H), 1.74-1.83 (m, 1H); ¹³C NMR (CD₃OD) δ 173.4, 171.7, 143.4, 140.1, 137.4, 134.0, 130.9, 129.7, 129.4, 125.5, 122.7, 53.0, 52.9, 31.3, 30.9, 15.1.

15 B. 4-[2(R)-tert-butoxycarbonylamino-3-triphenylmethylthiopropyl]amino-2-phenylbenzoyl-(S)-methionine methyl ester

To a mixture of 4-amino-2-phenylbenzoyl-(S)-methionine methyl ester hydrochloride (1.27 g, 3.22 mmol) in 20 mL of methanol was added N-Boc-S-trityl-(L)-cysteinyl (1.0 eq, according to ¹H NMR determination of aldehyde percentage) and sodium cyanoborohydride (400 mg, 2.0 eq). The mixture was stirred for 12 hr. After the evaporation of
20 solvents, the residue was extracted with ethyl acetate and concentrated sodium bicarbonate. After removing solvents, the residue was purified through flash column chromatography (1:1 = hexane: ethyl acetate, silica) to give the product 1.67 g
25 (yield 67%). ¹H NMR (CDCl₃) δ 7.65 (d, 8.6 Hz, 1H), 7.34-7.42 (m, 11H), 7.18-7.29 (m, 9H), 6.52 (dd, 2.3 and 8.1 Hz, 1H), 6.34 (d, 2.3 Hz, 1H), 5.65 (d, 7.7 Hz, 1H), 4.64 (ddd, 1H), 4.55 (d, 8.1 Hz, 1H), 4.19 (br t, 1H), 3.78 (br m, 1H), 3.64
30 (s, 3H), 3.09 (t, 6.1 Hz, 2H), 2.44 (m, 2H), 2.04-2.10 (m, 2H), 2.00 (s, 3H), 1.81-1.90 (m, 1H), 1.60-1.70 (m, 1H), 1.41 (s, 9H); ¹³C NMR (CDCl₃) δ

172.0, 168.3, 155.7, 149.4, 144.3, 141.6, 141.1,
131.3, 129.5, 128.7, 128.5, 127.9, 127.7, 126.8,
122.6, 113.6, 111.3, 79.8, 67.1, 52.2, 51.7, 49.5,
47.2, 34.3, 31.6, 29.4, 28.2, 15.2.

5 C. 4-[2(R)-amino-3-mercaptopropyl]amino-
2-phenylbenzoyl-(S)-methionine methyl
ester hydrochloride

The above N-Boc-S-trityl protected peptide
10 methyl ester (900 mg) was dissolved in 5 mL of
methanol. To this mixture was added a solution of
mercuric chloride (774 mg, 2.50 eq) in 5 mL of
methanol. The mixture was refluxed for 20 min.
The precipitate was collected and dried. This
15 solid was suspended in 10 mL of methanol and
reacted with gaseous hydrogen sulfide. After the
removal of black solid, the clear solution was
evaporated to dryness. The residue was then
dissolved in methylene chloride followed by
20 addition of 3N hydrogen chloride in ether. The
white solid was collected and dried to give the
pure product 476 mg (yield 81%). ¹H NMR (CD₃OD) δ
7.42 (d, 8.4 Hz, 1H), 7.30-7.38 (m, 5H), 7.77 (d,
8.4 Hz, 1H), 6.71 (s, 1H), 4.48 (dd, 4.2 and 5.1
25 Hz, 1H), 3.68 (s, 3H), 3.44-3.58 (m, 3H),
2.90-2.95 (dd, 4.1 and 14.5 Hz, 1H), 2.79-2.85
(dd, 4.7 and 14.5 Hz, 1H), 2.18-2.22 (m, 1H),
2.03-2.16 (m, 1H), 2.00 (s, 3H), 1.91-1.97 (m,
1H), 1.73-1.82 (m, 1H); ¹³C NMR (CD₃OD) δ 173.7,
30 173.4, 150.7, 143.5, 142.3, 131.2, 129.8, 129.5,
128.6, 125.6, 115.6, 112.2, 53.7, 53.2, 52.8,
45.0, 31.4, 30.9, 25.3, 15.0.

D. 4-[2(R)-amino-3-mercaptopropyl]amino-
2-phenylbenzoyl-(S)-methionine

35 The N-Boc-S-trityl protected peptide methyl
ester (500 mg) was hydrolyzed with 2.0 eq of
lithium hydroxide at 0°C for 1 hr. The product

was deprotected with trifluoroacetic acid (2 mL) in methylene chloride (1 mL). Triethylsilane was added dropwise until the deep yellow color disappeared. The mixture was kept at r.t. for 1.5
5 hr. After the evaporation of solvents, the residue was dried and washed with dry ether. The solid was purified through preparative HPLC to give a pure product 270 mg (yield 78%). ¹H NMR (CD₃OD) δ 7.44 (d, 8.4 Hz, 1H), 7.30-7.39 (m, 5H),
10 6.75 (d, 8.4 Hz, 1H), 6.67 (s, 1H), 4.45 (dd, 4.2 and 5.1 Hz, 1H), 3.42-3.58 (m, 3H), 2.90 (dd, 4.3 and 14.5 Hz, 1H), 2.81 (dd, 5.5 and 14.5 Hz, 1H),
2.17-2.23 (m, 1H), 2.09-2.15 (m, 1H), 2.00 (s, 3H), 1.90-1.99 (m, 1H), 1.71-1.81 (m, 1H); ¹³C NMR
15 (CD₃OD) δ 176.4, 173.5, 150.4, 143.0, 141.5, 131.0, 129.7, 129.4, 128.9, 124.6, 115.0, 112.3, 53.3, 49.6, 44.4, 30.8, 30.1, 24.9, 14.8.

E. 4-Nitro-2-phenylbenzoyl-(S)-leucine methyl ester

20 This compound was prepared through the coupling of 4-nitro-2-phenylbenzoic acid with (L)-leucine methyl ester hydrochloride as for the preparation of the methionine derivative (see Example 26, section A). ¹H NMR (CDCl₃) δ 8.24-8.26
25 (m, 2H), 7.86 (d, 8.7 Hz, 1H), 7.41-7.46 (m, 5H), 5.71 (d, 7.4 Hz, 1H), 4.57 (ddd, 1H), 3.67 (s, 3H), 1.37-1.46 (m, 1H), 1.08-1.25 (m, 2H), 0.78 (dd, 6H).

F. 4-[2(R)-tert-butoxycarbonyl-3-triphenylmethylthiopropyl]amino-2-phenylbenzoyl-(S)-leucine methyl ester

30 This compound was prepared using the same method as for the preparation of methionine derivative (See Example 26, section B), using 4-amino-2-phenylbenzoyl-(S)-leucine methyl ester and
35 N-Boc-S-trityl-(L)-cysteinyl as starting

materials. ¹H NMR (CDCl₃) δ 7.68 (d, 8.6 Hz, 1H), 7.33-7.41 (m, 11H), 7.17-7.29 (m, 9H), 6.50 (d, 8.6 Hz, 1H), 6.31 (s, 1H), 5.43 (d, 7.8 Hz, 1H), 4.60 (d, 6.1 Hz, 1H), 4.47 (ddd, 1H), 4.19 (br t, 1H), 3.77 (br m, 1H), 3.62 (s, 3H), 3.09 (t, 5.9 Hz, 2H), 2.45 (br m, 2H), 1.40 (s, 9H), 1.27-1.33 (m, 1H), 1.03-1.18 (m, 2H), 0.75 (dd, 6H); ¹³C (CDCl₃) δ 173.2, 168.2, 155.6, 149.4, 144.4, 141.7, 141.2, 131.4, 129.5, 128.8, 128.5, 127.9, 127.6, 126.8, 122.7, 113.6, 111.3, 79.6, 67.1, 51.9, 50.9, 49.5, 47.1, 41.2, 34.3, 28.3, 24.4, 22.7, 21.8.

G. 4-[2(R)-amino-3-mercaptopropyl]amino-2-phenylbenzoyl-(S)-leucine methyl ester hydrochloride

This compound was prepared with the same method as for the preparation of methionine derivative (see Example 26, section C), using 4-[2(R)-tert-butoxycarbonyl-3-triphenylmethylthiopropyl]amino-2-phenylbenzoyl-(S)-leucine methyl ester and mercuric chloride. ¹H NMR (CD₃OD) δ 7.42 (d, 8.5 Hz, 1H), 7.31-7.38 (m, 5H), 6.76 (d, 8.5 Hz, 1H), 6.68 (s, 1H), 4.33 (t, 7.8 Hz, 1H), 3.67 (s, 3H), 3.46-3.55 (m, 3H), 2.95 (dd, 4.4 and 14.5 Hz, 1H), 2.81 (dd, 5.1 and 14.5 Hz, 1H), 1.44 (t, 7.6 Hz, 2H), 1.18-1.25 (m, 1H), 0.76-0.83 (dd, 4.1 and 6.6 Hz, 6H).

H. 4-[2(R)-amino-3-mercaptopropyl]amino-2-phenylbenzoyl-(S)-leucine

This compound was prepared with the same method as for the preparation of the methionine derivative (see Example 26, section D), using 4-[2(R)-amino-3-mercaptopropyl]amino-2-phenylbenzoyl-(S)-leucine methyl ester hydrochloride and lithium hydroxide. ¹H NMR (CD₃OD) δ 7.42 (d, 8.5 Hz, 1H), 7.29-7.38 (m, 5H),

6.73 (d, 8.5 Hz, 1H), 6.66 (s, 1H), 4.32 (dd, 3.3
and 5.9 Hz, 1H), 3.41-3.57 (m, 3H), 2.94 (dd, 4.3
and 14.5 Hz, 1H), 2.78 (5.2 and 14.5 Hz, 1H), 1.45
(t, 6.7 Hz, 2H), 1.17-1.26 (m, 1H), 0.78-0.83 (t,
5 8.5 Hz, 6H).

I. 4-Nitro-2-naphthylbenzoic acid

The coupling of 4-nitro-2-bromobenzoic acid
methyl ester (1.92 g, 7.4 mmol) with 1-
naphthylboronic acid (2.53 g, 14.7 mmol) in the
10 presence of anhydrous sodium phosphate (3.64 g,
22.2 mmol) and palladium
tetrakis(triphenylphosphine) (426 mg, 0.368 mmol) in
50 mL of DMF at 100°C gave the
4-nitro-2-naphthylbenzoic acid methyl ester (1.66
15 g, 73% yield). ¹H NMR (CDCl₃) δ 8.34 (d, 8.5 Hz,
1H), 8.28 (s, 1H), 8.14 (d, 8.5 Hz, 1H), 7.92 (d,
8.2 Hz, 2H), 7.47-7.56 (m, 2H), 7.41 (d, 3.8 Hz,
2H), 7.34 (d, 6.8 Hz, 1H), 3.40 (s, 3H). After
the hydrolysis of methyl ester, 1.44 g of product
20 was collected (yield 91%). ¹H NMR (CDCl₃) δ 8.33
(d, 8.6 Hz, 1H), 8.23 (s, 1H), 8.17 (d, 8.6 Hz,
1H), 7.88 (d, 8.2 Hz, 2H), 7.46-7.52 (m, 2H),
7.38-7.42 (m, 2H), 7.33 (d, 7.0 Hz, 1H); ¹³C NMR
(CD₃COCD₃) δ 167.2, 150.1, 143.1, 139.0, 138.2,
25 134.4, 132.5, 132.1, 129.2, 127.3, 126.7, 125.8,
125.9, 123.3.

**J. 4-Nitro-2-naphthylbenzoyl-
(S)-methionine methyl ester**

The coupling of 4-nitro-2-naphthylbenzoic
30 acid with (L)-methionine methyl ester in the
presence of EDCI and HOBT provided the desired
product (yield 95%). TLC of the product showed
single spot, but ¹H NMR showed the presence of
diastereomers caused by the restricted rotation
35 between naphthyl and phenyl rings. ¹H NMR (CDCl₃)
δ 8.33-8.38 (m, 1H), 8.26 (ss, 1H), 8.14 (d, 8.5

Hz, 0.5H), 8.00 (d, 8.5 Hz, 0.5H), 7.94-7.98 (m, 2H), 7.42-7.65 (m, 5H), 5.98 (t, 1H), 4.42 (m, 1H), 3.56 (s, 1.5H), 3.51 (s, 1.5H), 1.83 (s, 1.5H), 1.74 (s, 1.5H), 1.56-1.64 (m, 1H),
5 1.33-1.45 (m, 2H), 1.09-1.14 (m, 1H).

K. 4-[2(R)-tert-butoxycarbonyl-3-triphenylmethylthiopropyl]amino-2-naphthyl-(S)-methionine methyl ester

The reduction of

10 4-nitro-2-naphthylbenzoyl-(L)-methionine methyl ester gave a quantitative yield of the amino derivative which was reacted with N-Boc-S-trityl cysteinal in the presence of sodium
15 cyanoborohydride. After flash column chromatography (1:1 = ethyl acetate : hexane) purification, a desired product was obtained (yield 40%). TLC showed single spot, but ¹H NMR
20 showed diastereomers caused by the restricted rotation between the naphthyl and phenyl rings. ¹H NMR (CDCl₃) δ 7.84-7.96 (m, 3H), 7.49-7.66 (m, 4H), 7.37-7.43 (m, 7H), 7.14-7.27 (m, 9H), 6.60-6.63 (d, 8.6 Hz, 1H), 6.33 (m, 1H), 5.67 (d, 7.8 Hz, 0.6H), 5.60 (d, 7.8 Hz, 0.4H), 4.56 (br d, 6.2 Hz, 1H), 4.35-4.44 (m, 1H), 4.30 (br, 1H), 3.78 (br m, 1H), 3.55 (s, 1.9H), 3.38 (s, 1.1H), 3.06 (t, 5.8 Hz, 2H), 2.44 (m, 2H), 1.90 (s, 1H), 1.79 (s, 2H), 1.57-1.68 (m, 0.5H), 1.36-1.45 (m, 10H), 1.23-1.32 (m, 2H), 0.94-0.98 (m, 0.7H).

L. 4-[2(R)-amino-3-mercaptopropyl]amino-2-naphthylbenzoyl-(S)-methionine

30 This compound was prepared from the N-Boc-S-trityl protected form (section K) by saponification followed with acidic cleavage by trifluoroacetic acid. The pure compound was
35 obtained through preparative HPLC. ¹H NMR showed complicated diastereomers caused by the restricted

rotation of aryl-aryl bond. ^1H NMR (CD_3OD) δ
7.86-7.94 (m, 2H), 7.73 (d, 8.6 Hz, 0.6H),
7.35-7.67 (m, 5.4H), 6.83-6.88 (m, 1H), 6.63-6.67
(m, 1H), 4.17-4.23 (m, 1H), 3.41-3.58 (m, 3H),
5 2.91 (dd, 4.2 and 14.5 Hz, 1H), 2.80 (dd, 5.3 and
14.5 Hz, 1H), 1.82 (s, 1.4H), 1.80 (s, 1.6H),
1.65-1.77 (m, 1H), 1.41-1.52 (m, 2H), 1.09-1.32
(m, 1H).

10 M. 4-Nitro-2-naphthylbenzoyl-
(S)-leucine methyl ester

This compound was prepared with the same
method as for the preparation of the methionine
derivative (section J) using 4-nitro-2-
naphthylbenzoic acid, (S)-leucine methyl ester,
15 EDCI and HOBT. ^1H NMR (CDCl_3) δ 8.34-8.39 (m, 1H),
8.25 (s, 1H), 8.18 (d, 8.6 Hz, 0.6H), 8.02 (d, 8.6
Hz, 0.4H), 7.91-8.00 (m, 2H), 7.62 (t, 7.0 Hz,
0.6H), 7.48-7.58 (m, 3H), 7.41 (t, 7.0 Hz, 1.4H),
5.71 (d, 7.9 Hz, 0.6H), 5.60 (d, 7.9 Hz, 0.4H),
20 4.29 (m, 1H), 3.57 (s, 1.7H), 3.52 (s, 1.3H),
1.05-1.11 (m, 0.5H), 0.88-0.97 (m, 0.7H),
0.69-0.78 (m, 0.5H), 0.41-0.59 (m, 7.0H),
0.19-0.26 (m, 0.6H).

25 N. 4-[2(R)-tert-butoxycarbonylamino-3-
triphenylmethylthiopropyl]amino-2-
naphthylbenzoyl-(S)-leucine methyl ester

This compound was prepared with the same
method as for the preparation of the methionine
derivative (section K). ^1H NMR (CDCl_3) δ 7.85-8.00
30 (m, 3H), 7.47-7.67 (m, 4H), 7.39-7.43 (m, 7H),
7.14-7.37 (m, 9H), 6.61 (d, 8.6 Hz, 1H), 6.32 (s,
1H), 5.46 (d, 7.6 Hz, 0.6H), 5.36 (d, 7.6 Hz,
0.4H), 4.55 (d, 7.2 Hz, 1H), 4.20-4.27 (m, 2H),
3.76 (br, 1H), 3.56 (s, 2H), 3.38 (s, 1H), 3.06
35 (t, 5.9 Hz, 2H), 2.43 (m, 2H), 1.36-1.43 (m, 9H),

0.81-1.03 (m, 1H), 0.55-0.67 (m, 2.8H), 0.36-0.45 (m, 4.7H), 0.00-0.09 (m, 0.6H).

5 **O. 4-[2(R)amino-3-mercaptopropyl]amino-2-naphthylbenzoyl-(S)-leucine methyl ester**

 This compound was prepared from the N-Boc-S-trityl methyl ester of the corresponding compound, using the method of section L. ¹H NMR (CDCl₃) δ 7.98 (d, 8.5 Hz, 0.6H), 7.84-7.90 (m, 2.4H), 7.65 (d, 8.5 Hz, 0.4H), 7.43-7.58 (m, 3.6H), 7.34-7.39 (m, 1H), 6.72 (m, 1H), 6.45 (ss, 1H), 5.46 (d, 7.8 Hz, 0.6H), 5.40 (d, 7.7 Hz, 0.4H), 4.64 (m, 1H), 4.23 (m, 1H), 3.54 (s, 2H), 3.30 (s, 1H), 3.25 (m, 1H), 2.97-3.06 (m, 2H), 2.67 (dd, 3.7 and 13.1 Hz, 1H), 2.47 (dd, 6.5 and 13.2 Hz, 1H), 1.45-1.65 (br s, 2H), 0.81-1.03 (m, 1.2H), 0.54-0.67 (m, 3H), 0.36-0.39 (m, 4.3H), 0.00-0.10 (m, 0.7H).

EXAMPLE 27

20 **FTase and GGTase I Activity Assay**

 FTase and GGTase I activities from 60,000 x g supernatants of human Burkitt lymphoma (Daudi) cells (ATCC, Rockville, MD) were assayed exactly as described previously for FTase (41).

25 Inhibition studies were performed by determining the ability of Ras CAAX peptidomimetics to inhibit the transfer of [³H]-farnesyl and [³H]-geranylgeranyl from [³H]FPP and [³H]GGPP to H-ras-CVLS and H-Ras-CVLL, respectively (41).

30

EXAMPLE 28

Ras and Rap1A Processing Assay

 H-Ras cells (45) and K-Ras4B Cells (32) were kind gifts from Dr. Channing Der and Dr. Adrienne Cox (University of North Carolina, Chapel Hill).

Means of obtaining these cell lines will be easily recognized by the skilled practitioner.

Cells were seeded on day 0 in 100 mm dishes in Dulbecco's modified Eagles medium supplemented with 10% calf serum and 1% penicillin-streptomycin. On days 1 and 2, cells were refed with medium containing various concentrations of FTI-277, GGTI-286 or vehicle (10 mM DTT in DMSO). On day 3, cells were washed and lysed in lysis buffer containing 50 mM HEPES, pH 7.5, 10 mM NaCl, 1% TX-100, 10% glycerol, 5 mM MgCl₂, 1 mM EGTA, 25 µg/ml leupeptin, 2 mM PMSF, 2 mM Na₃VO₄, 1 mg/ml soybean trypsin inhibitor, 10 µg/ml aprotinin, 6.4 mg/ml Sigma-104[®] phosphatase substrate. Lysates were cleared (14,000 rpm, 4°C, 15 min) and equal amounts of protein were separated on a 12.5% SDS-PAGE, transferred to nitrocellulose, and immunoblotted using an anti-Ras antibody (Y13-259, ATCC) or an anti-Rap1A antibody (SC-65, Santa Cruz Biotechnology, Santa Cruz, CA). Antibody reactions were visualized using either peroxidase-conjugated goat anti-rat IgG (for Y13-259), or peroxidase-conjugated goat anti-rabbit IgG (for Rap1A) and an enhanced chemiluminescence detection (ECL, Amersham Corp.), as described previously (41).

EXAMPLE 29

MAP Kinase Immunoblotting

Cells were treated with FTI-277, GGTI-286, or vehicle and lysed as previously described for Ras and Rap1A processing. Equal amounts of protein were separated on a 15% SDS-PAGE, transferred to nitrocellulose, and immunoblotted using an anti-MAP kinase antibody (erk2, monoclonal, UB1, Lake Placid, NY). Antibody reactions were visualized using peroxidase-conjugated donkey anti-mouse IgG

(Jackson ImmunoResearch Laboratories Inc., West Grove, PA) and an enhanced chemiluminescence detection system (ECL, Amersham Corp.)

EXAMPLE 30

5 ***Inhibition of GGTase I by GGTI-286***

GGTI-287 potentially inhibited GGTase I *in vitro* (IC_{50} =5 nM) and was selective towards inhibiting GGTase I over FTase (IC_{50} =25 nM) (Table 5). Thus, the substitution of methionine in FTI-276 by a
10 leucine in GGTI-287 (Fig. 17) increased the potency towards GGTase I by approximately 10-fold (Table 5). More importantly, it reversed the selectivity from a FTase to a GGTase I-specific inhibitor by a factor of 500 (Table 5). To
15 determine whether this selectivity is respected in whole cells, the cell-permeable methyl ester derivative of GGTI-287, GGTI-286 (Fig. 17), was synthesized and used to treat NIH 3T3 cells which overexpress oncogenic H-Ras-CVLS (31). Cell
20 lysates were electrophoresed on SDS-PAGE and immunoblotted with an anti-Ras antibody as described in Example 28. Figure 18 shows that accumulation of unprocessed H-Ras did not occur at concentrations lower than 30 μ M GGTI-286.
25 Therefore, GGTI-286 is not a good inhibitor of H-Ras processing in whole cells. However, GGTI-286 was a very potent inhibitor of the processing of the geranylgeranylated Rap1A protein (IC_{50} =2 μ M) (Fig. 18). Thus, GGTI-286 is more than 15-fold
30 selective for inhibition of geranylgeranylation over farnesylation processing (Table 5). This data is in direct contrast to the FTase specific inhibitor FTI-277 which inhibited H-Ras and Rap1A processing with IC_{50} s of 100 nM and 50 μ M,
35 respectively (Fig. 18). Thus, GGTI-286 is 25-fold

more potent than FTI-277 at inhibiting geranylgeranylation in whole cells (Table 5).

EXAMPLE 31

Inhibition of GGTase I by GGTI-297 and GGTI-298

5 To determine the effect of replacing the phenyl substituent with a naphthyl on GGTase I inhibition, 4-[2(R)-amino-3-mercaptopropyl] amino-2-naphthyl benzoyl-(L)-leucine (GGTI-297) and its methylester (GGTI-298) were tested. GGTI-297
10 inhibited GGTase I in vitro with an IC_{50} of 40 nM and was selective towards inhibiting GGTase I over FTase (IC_{50} = 270 nM) (Fig. 21, Table 5). Thus, the substitution of phenyl in GGTI-287 by naphthyl in GGTI-297 decreased the potency towards GGTase I by
15 8 fold and towards FTase by over 10-fold. However, more importantly, the selectivity for GGTase I over FTase increased from 5-fold (GGTI-287) to 7-fold (GGTI-297), as shown in Table 5. This selectivity was also respected in vivo since
20 concentrations as high as 20 μ M did not inhibit H-Ras processing whereas Rap1A and Ras4B were completely blocked at 10 μ M GGTI-298 (Table 5).

EXAMPLE 32

Inhibition of K-Ras4B Function by GGTI-286

25 The ability of GGTI-286 to inhibit the processing and signaling of oncogenic K-Ras4B was then evaluated. NIH 3T3 cells which overexpress oncogenic K-Ras4B (32) were treated either GGTI-286 (0-30 μ M) or FTI-277 (0-30 μ M) and the lysates
30 were immunoblotted with an anti-Ras antibody as described under Example 28. Figure 19 shows that GGTI-286 inhibited potently the processing of K-Ras4B with an IC_{50} of 2 μ M. The ability of GGTI-286 to inhibit the processing of K-Ras4B was much
35 closer to its ability to inhibit the processing of

geranylgeranylated Rap1A ($IC_{50}=2 \mu M$) than that of
farnesylated H-Ras ($IC_{50}>30 \mu M$) (Fig. 18) (Table 5).
This suggested that K-Ras4B might be
geranylgeranylated. Consistent with this is the
5 fact that K-Ras4B processing was very resistant to
the FTase-specific inhibitor FTI-277 ($IC_{50}=10 \mu M$)
(Fig. 19). Furthermore, GGTI-286 inhibited K-
Ras4B processing at concentrations (1-3 μM) (Fig.
19) that had no effect on the processing of
10 farnesylated H-Ras (Fig. 18).

EXAMPLE 33

Effects of GGTI-286 on Oncogenic K-Ras 4B Constitutive Activation of MAP Kinase

To determine whether inhibition of K-Ras4B
15 processing by GGTI-286 results in disruption of
oncogenic signaling, the ability of GGTI-286 to
antagonize oncogenic K-Ras 4B constitutive
activation of MAP kinase was examined. Activated
MAP kinase is hyperphosphorylated and migrates
20 slower than hypophosphorylated (inactive) MAP
kinase on SDS-PAGE (43, 66). Figure 20 shows that
K-Ras4B transformed cells contained mainly
activated MAP kinase. Treatment of these cells
with the FTase-specific inhibitor FTI-277 (0-30
25 μM) did not inhibit MAP kinase activation until 30
 μM (Fig. 20). In contrast, GGTI-286 inhibited MAP
kinase activation with an IC_{50} of 1 μM and the
block was complete at 10 μM . Thus, GGTI-286
blocked oncogenic K-Ras4B MAP kinase activation at
30 a concentration (10 μM) where FTI-277 had no
effect. In contrast, oncogenic H-Ras activation
of MAP kinase was inhibited only slightly by GGTI-
286 whereas FTI-277 completely blocked this
activation at 3 μM (Fig. 20). Furthermore, GGTI-
35 286 blocked K-Ras4B activation of MAP kinase at a

concentration (10 μ M) that had little effect on H-Ras activation of MAP kinase (Fig. 20).

EXAMPLE 34

Antitumor Efficacy

5 The above examples demonstrate that GGTI-286 is a potent and highly selective inhibitor of K-Ras4B processing and activation of oncogenic signalling. In order to demonstrate the efficacy of these inhibitors as anticancer agents, K-Ras4B
10 transformed NIH-3T3 cells were implanted subcutaneously in nude mice. When the tumors reached sizes of 50-100 mm³, the mice were randomly separated into control and treated groups (5 animals per group, each animal had a tumor on both
15 the right and the left flank). Figure 22 shows that tumors from control animals treated with saline once daily grew to an average size of 2900 mm³ over a period of two weeks. In contrast, tumors from animals treated once daily with GGTI-
20 286 (25 mg/kg or 50 mg/kg) grew to a size of 1600 mm³ or 900 mm³, respectively (Fig. 22). Thus, GGTI-286 inhibited tumor growth by 50% and 70%, respectively.

25 In summary, the data clearly identifies GGTI-286 not only as a potent antagonist of K-Ras4B oncogenic signaling in cultured cells, but also as an inhibitor of tumor growth in whole animals.

30 References cited herein are listed below for convenience and are hereby incorporated by reference.

REFERENCES

1. Barbacid, M., Annu. Rev. Biochem.,
56:779-828, (1987)
- 5 2. Grand, R.J.A. and Owen, D., Biochem. J.,
279:609-631, (1991)
3. Barbacid, M., Important Advances in
Oncology, eds. Devita, Hellman &
Rosenberg (Lippincott, Philadelphia,
PA), pp. 3-22, (1986)
- 10 4. Mulcahy, L.S., Smith, M.R., and Stacey,
D.W., Nature, 313:241-243, (1985)
5. Noda, M., Ko, M., Ogura, A., Liu, D.G.,
Amano, T., Takano, T. and Ikawa, Y.,
Nature, 318:73-75, (1985)
- 15 6. Bar-Sagi, D. and Feramisco, J.R., Cell,
42:841-848, (1985)
7. Kataoka, T., Powers, S., McGill, C.,
Fasano, O., Strathern, J., Broach, J.
and Wigler, M., Cell, 37:437-445, (1984)
- 20 8. Willumsen, B.M., Christensen, A.,
Hubbert, N.C., Papageorge, A.G. and
Lowy, D.R., Nature, 310:583-586, (1984)
- 25 9. Willumsen, B.M., Norris, K., Papageorge,
A.G., Hubbert, N.C. and Lowy, D.R., EMBO
J., 3:2581-2585, (1984)
10. Hancock, J.F., Magee, A.I., Childs, J.E.
and Marshall, C.J., Cell, 57:1167-1177,
(1989)
- 30 11. Gutierrez, L., Magee, A.I., Marshall,
C.J. and Hancock, J.F., EMBO J., 8:1093-
1098, (1989)
12. Casey, P.J., Soliski, P.A., Der, C.J. and
Buss, J.E., Proc. Natl. Acad. Sci. USA,
86:8323-8327, (1989)
- 35 13. Jackson, J.H., Cochrane, C.G., Bourne,
J.R., Soliski, P.A., Buss, J.E. and Der,
C.J., Proc. Natl. Acad. Sci. USA,
87:3042-3046, (1990)
- 40 14. Hancock, J.F., Paterson, H. and
Marshall, J.C., Cell, 63:133-139, (1990)

15. Reiss, Y., Goldstein, J.L., Seabra, M.C., Casey, P.J. and Brown, M.S., Cell, 62:81-88, (1990)
- 5 16. Reiss, Y., Seabra, M.C., Goldstein, J.L. and Brown, M.S., METHODS: A Companion to Methods in Enzymology, 1:241-245, (1990)
- 10 17. Reiss, Y., Seabra, M.C., Armstrong, S.A., Slaughter, C.A., Goldstein, J.L. and Brown, M.S., J. Biol. Chem., 266:10672-10877, (1991)
- 15 18. Reiss, Y., Stradley, S.J., Gierasch, L.M., Brown, M.S. and Goldstein, J.L., Proc. Natl. Acad. Sci. USA, 88:732-736, (1991)
- 20 19. Manne, V., Roberts, D., Tobin, A., O'Rourke, E., DeVirgillio, M., Meyres, C., Ahmed, N., Kurz, E., Resh, M., Kung, H.F. and Barbacid, M., Proc. Natl. Acad. Sci. USA, 87:7541-7545, (1990)
- 20 20. Gibbs, J.B., Cell, 65:1-4, (1991)
- 25 21. Moores, S.L., Schaber, M.D., Mosser, S.D., Rands, E., O'Hara, M.B., Garsky, V.M., Marshall, M.S., Pompliano, D.L. and Gibbs, J.B., J. Biol. Chem., 266:14603-14610, (1991)
- 30 22. Goldstein, J.L., Brown, M.S., Stradley, S.J., Reiss, Y. and Gierasch, L.M., J. Biol. Chem., 266:15575-15578, (1991)
- 30 23. Brown, M.S., Goldstein, J.L., Paris, K.J., Burnier, J.P. and Marsters, J.C., Proc. Natl. Acad. Sci. USA, 89:8313-8316, (1992)
- 35 24. Pompliano, D.L., Rands, E., Schaber, M.D., Mosser, S.D., Neville, J.A. and Gibbs, J.B., Biochemistry, 31:3800-3807, (1992)
- 40 25. Lacal, J.D., Santos, E., Notario, V., Barbacid, M., Yamazaki, S., Kung, H.F., Seamans, O., McAndrew, S. and Crowl, R., Proc. Natl. Acad. Sci. USA, 81:5305-5309, (1984)
26. Stewart, F.H.C., Aus. J. Chem., 36:2511, (1983)

27. Brown, M.J., Milano, P.D., Lever, D.C.,
Epstein, W.W. and Poulter, C.D., J. Am.
Chem. Soc., 113:3176, (1991)
- 5 28. Yang, C.C., Marlowe, C.K. and Kania, R.,
J. Am. Chem. Soc., 113:3177, (1991)
29. McCormick, F. (1993) Nature 363:15-
16.
30. McCormick, F. (1994) Current Opinion in
Genetics & Development 4:71-76
- 10 31. Marshall, C.J. (1994) Current Opinion in
Genetics & Development 4: 82-89
32. Kato, K., Cox, A.D., Hisaka, M.M.,
Graham, S.M., Buss, J.E., and Der, C.J.
15 (1992) Proc. Natl. Acad. Sci. USA
89:6403-6407.
33. Gibbs, J.B., Oliff, A., and Kohl, N.E.
(1994) Cell 77:175-178.
34. Nigam, M., Seong, C., Qian, Y.,
20 Hamilton, A.D., and Sebti, S. M. (1993)
J. Biol. Chem. 268:20695-20698.
35. Qian, Y., Blaskovich, M.A., Saleem, M.,
Seong, C., Wathen, S.P., Hamilton, A.D.,
and Sebti, S.M. (1994) J. Biol. Chem.
269:12410-12413.
- 25 36. Qian, Y., Blaskovich, M.A., Seong, C.M.,
Vogt, A., Hamilton, A.D., and Sebti,
S.M. (1994) Bioorg. Med. Chem. Lett
4:2579-2584.
- 30 37. Kohl, N.E., Mosser, S.D., deSolms, S.J.,
Giuliani, E.A., Pompliano, D.L., Graham,
S.L., Smith, R.L., Scolnick, E.M.,
Oliff, A., and Gibbs, J.B. (1993)
Science 260:1934-1937.
- 35 38. James, G.L., Goldstein, J.L., Brown,
M.S., Rawson, T.E., Somers, T.C.,
McDowell, R.S., Crowley, C.W., Lucas,
B.K., Levinson, A.D., and Marsters,
J.C., Jr. (1993) Science 260:193-194.
- 40 39. Graham, S.L., deSolms, S.J., Giuliani,
E.A., Kohl, N.E., Mosser, S.D., Oliff,
A.I., Pompliano, D.L., Rands, E.
Breslin, M.J., Deana, A.A., Garsky,
V.M., Scholz, T.H., Gibbs, J.B., and

- Smith, R.L. (1994) J. Med. Chem. 37:725-732.
- 5 40. Garcia, A.M., Rowell, C. Ackerman, K., Kowalczyk, J.J., and Lewis, M.D. (1993) J. Biol. Chem. 268:18415-18418.
41. Vogt, A., Qian, Y., Blaskovich, M.A., Fossum, R.D., Hamilton, A.D., and Sebt, S.M. (1995) J. Biol. Chem. 270:660-664.
- 10 42. Kohl, N.E., Wilson, F.R., Mosser, S.D., Giuliani, E., deSolms, S.J., Conner, M.W., Anthony, N.J., Holtz, W.J., Gomez, R.P., Lee, T.J., and et al. (1994) Proc. Natl. Acad. Sci. USA 91:9141-9145.
- 15 43. Cox, A.D., Garcia, A.M., Westwick, J.K., Kowalczyk, J.J., Lewis, M.D., Brenner, D.A., and Der, C.J. (1994) J. Biol. Chem. 269:19203-19206.
- 20 44. James, G.L., Brown, M.S., Cobb, M.H., and Goldstein, J.L. (1994) J. Biol. Chem. 269:277705-277714.
45. Cox, A.D., Hisaka, M.M., Buss, J.E., and Der, C.J. (1992) Mol. Cell. Biol. 12:2606-2615.
- 25 46. Hallberg, B. Rayter, S.I., and Downward, J. (1994) J. Biol. Chem. 269:3913-3916.
47. Leever, S.J., Paterson, H.F., and Marshall, C.J. (1994) Nature 369:411-414.
- 30 48. Stanton, V. P., Jr., Nichols, D. W., Laudano, A. P., Cooper, G. M. (1989) Mol. Cell. Biol. 9: 639-647.
49. Casey, P., J. Lipid. Res., 88:1731-1740 (1992)
- 35 50. Cox et al, Curr. Op. Cell Biol., 4:1008-1016 (1992)
51. Goldstein et al, Science, 260:1937-1942 (1993)
- 40 52. Watanabe et al, Syn. Lett., 3:207-210 (1992)

53. Stradley et al, Biochemistry, 32:12586-12590 (1993)
54. Gardino, J. et al, J. Org. Chem., 49: 5237-5243 (1984)
- 5 55. Wallow, T. I. et al. J. Org. Chem.: 59, 5034-5037 (1994)
56. Crowther, G. P. et al., Org. Syn., 51: 96-100 (1971)
- 10 57. Fincham, C. I. et al., J. Med. Chem., 35: 1472-1484 (1992)
58. Goel, O. P. et al., Org. Syn., 67: 69-75 (1989)
59. Pearson, D. A. et al., Tetrahedron Lett., 30: 2739-2742 (1989)
- 15 60. Casey, P.J., Solski, P.A., Der, C.J., and Buss, J.E. (1989) *Cell* 57, 1167-1177
61. Seabra, M.C., Reiss, Y., Casey, P.J., Brown, M.S., and Goldstein, J.L. (1991) *Cell* 65, 429-434
- 20 62. Zhang, F.L., Diehl, R.E., Kohl, N.e., Gibbs, J.B., Giros, B., Casey, P.J., and Omer, C.A. 91994) *J. Biol. Chem.* 269, 3175-3180
- 25 63. Omer, C.A., Kral, A.M., Diehl, R.E., Prendergast, G.C., Powers, S., Allen, C.M., Gibbs, J.B., and Kohl, N.E. (1993) *Biochemistry* 32, 5167-5176
- 30 64. Yokoyama, K., Goodwin, G.W., Ghomashchi, F., Glomser, J.A., and Gelb, M.H. (1991) *Proc. Natl. Acad. Sci. USA* 88, 5302-5306
65. Trueblood, C.E., Ohya, Y., and Rine, J. (1993) *Mol. Cell. Biol.* 13, 4260-4275
- 35 66. Lerner, E.G., Qian, Y., Blaskovich, M.A., Fossum, R.D., Vogt A., Sun, J., Cox, A.D., Der, C.J., Hamilton, A.D., and Sebti, S.M. (1995) *J. Biol. Chem.* 270, 26802-26806
- 40 67. James, G.L., Goldstein, L.L., and Brown, M.S. (1995) *J. Biol. Chem.* 270, 6221-6226

68. Sun, J., Qian, Y., Hamilton, A.D., and
Sebti, S.M. (1995) *Cancer Research* 55,
4243-4247.

69. Moomaw, J.P. and Casey, P.J. (1992) *J.*
Biol. Chem. 267, 17438-17443

5

It will be appreciated that various

modifications may be made in the invention as
described above without departing from the scope
and intent of the invention as defined in the

10

following claims wherein:

CLAIMS

WE CLAIM:

1. A peptidomimetic of the formula:

$C\beta X$

5 wherein

C is a cysteinyl moiety, in a reduced or nonreduced state;


X is an amino acid; and

10 β is a residue of an aminobenzoic acid or an aminonaphthoic acid.

2. A peptidomimetic according to claim 1 wherein the cysteinyl moiety is in the reduced state.

15 3. A peptidomimetic according to claim 2 wherein β is 2-phenyl-4-aminobenzoic acid.

4. A peptidomimetic according to claim 1 of the formula:



20 5. A peptidomimetic according to claim 1 wherein β is a substituted 4-aminobenzoic acid.

6. A peptidomimetic according to claim 1 wherein X is methionine or phenylalanine.

7. A pharmaceutical composition comprising a peptidomimetic according to claim 1 and a pharmaceutically acceptable carrier.

5 8. A method of inhibiting farnesyltransferase in a host where the farnesyltransferase is present which comprises administering to the host an effective amount of a peptidomimetic according to claim 1.


10 9. A peptidomimetic according to claim 1 in the form of a pro-drug.

15 10. A pro-drug according to claim 9 wherein the pro-drug comprises a compound as defined with one or more terminal amino, sulfhydryl and acid groups functionalized with a lipophilic, esterase-sensitive moiety.

20 11. A pro-drug according to claim 9 wherein the terminal amino and sulfhydryl groups of the cysteine C are functionalized by benzyloxy carbonyl groups or other cleavable lipophilic groups and a terminal carboxylic acid group is esterified.

12. A pro-drug according to claim 11 wherein the carboxylic acid group is esterified as the methyl ester.

13. A compound of the formula:




wherein R represents H, CH₃, or any
lipophilic esterase-sensitive moiety,
and R¹ represents H or a substituted or
unsubstituted phenyl group.

14. A compound according to claim 13 wherein
R¹ is an unsubstituted phenyl group, or an alkoxy-,
chloro-, bromo- or methyl- substituted phenyl
group.

15. A compound according to claim 13 wherein
R¹ is chosen from the group consisting of a 3,5
dimethylphenyl radical, a thiophene radical, a
naphthyl radical, a pyrrole radical, a pyridyl
radical, an alkyl radical, and an alkoxy radical.

16. A compound of the formula




wherein R represents H, CH₃, or any lipophilic esterase-sensitive moiety,

and R¹ represents H or a substituted or unsubstituted phenyl group.

- 5 17. A compound according to claim 15 wherein R¹ is an unsubstituted phenyl group, or an alkoxy-, chloro-, bromo- or methyl- substituted phenyl group.


- 10 18. A compound according to claim 16 wherein R¹ is a 3,5 dimethylphenyl radical.

19. A compound of the formula



wherein R represents H, CH₃, or any lipophilic esterase-sensitive moiety.

20. A compound of the formula



wherein R represents H, CH₃, or any lipophilic
esterase-sensitive moiety,
and R¹ represents H, CH₃, or OCH₃.

21. A method of inhibiting
5 farnesyltransferase in a host wherein the
farnesyltransferase is present which comprises
administering to the host an effective amount of a
peptidomimetic according to claim 13.
22. A method of treating cancer comprising
10 administering to a patient in need of such
treatment an effective amount of a peptidomimetic
according to claim 1 or 13.
23. A peptidomimetic according to claim 1
wherein β is a residue of an aminobenzoic acid.
24. A peptidomimetic according to claim 23
15 wherein X is leucine or isoleucine.
25. A peptidomimetic according to claim 24
wherein β is a substituted 4-aminobenzoic acid.
26. A peptidomimetic according to claim 25
20 wherein the cysteinyl moiety is in the reduced
state.
27. A peptidomimetic according to claim 26
wherein the 4-aminobenzoic acid is substituted at
the 2- and/or 3- position of the phenyl ring by an
25 alkyl, alkoxy, aryl, or naphthyl group, or a
heterocyclic or heteroaromatic ring.
28. A peptidomimetic according to claim 26
wherein β is 2-phenyl-4-aminobenzoic acid or 2-
naphthyl-4-aminobenzoic acid.

29. A peptidomimetic according to claim 28 wherein L is leucine.

5 30. A pharmaceutical composition comprising a peptidomimetic according to claim 24 and a pharmaceutically acceptable carrier.

10 31. A method of inhibiting geranylgeranyltransferase in a host where the geranylgeranyltransferase is present which comprises administering to the host an effective amount of a peptidomimetic according to claim 24.

32. A peptidomimetic according to claim 24 in the form of a pro-drug.

15 33. A pro-drug according to claim 32 wherein the pro-drug comprises a compound as defined with one or more terminal amino, sulfhydryl and acid groups functionalized with a lipophilic, esterase-sensitive moiety.

34. A pro-drug according to claim 33 wherein a terminal carboxylic acid group is esterified.

20 35. A pro-drug according to claim 34 wherein the carboxylic acid group is esterified as the methyl ester.

25 36. A method of inhibiting geranylgeranyltransferase in a host where geranylgeranyltransferase is present which comprises administering to the host an effective amount of the compound of claim 35.

37. A method of treating cancer comprising administering to a patient in need of such

treatment an effective amount of a peptidomimetic according to claim 24.


38. A method of treating cancer comprising administering to a patient in need of such treatment an effective amount of a peptidomimetic according to claim 35.

39. A compound of the formula



wherein

10



15

A representing O or 2H, and

R_0 representing SH, NH_2 , or $C_xH_y-SO_2-NH-$,

wherein C_xH_y is a straight chain saturated or unsaturated hydrocarbon, with x being between 1 and 20 and y between 3 and 41, inclusive; and

20

B stands for $-NHR$, wherein R is an aryl group.

40. A compound according to claim 39 wherein R is a biphenyl group.

41. A compound according to claim 39 wherein C^0 is 3-mercapto-2-amino-propylamino.

25


42. A compound according to claim 41 wherein R is a biphenyl group.

43. A compound according to claim 42 wherein R is a biphenyl group substituted with $-COOH$.

44. A compound according to claim 42 wherein R is a biphenyl group substituted with lower alkyl or oxyalkyl.

45. A compound according to claim 44 wherein the lower alkyl is methyl.

46. A compound according to claim 39 of the formula:



wherein A is O or 2H; R₁ is hydrogen or COOH; R₂ is hydrogen, COOH, COOCH₃, CH₃, or tetrazolyl; R₃ is hydrogen or COOH; and R₄ is hydrogen, OCH₃, OC₂H₅, or a phenyl group.

47. A compound according to claim 46 wherein R₁ is hydrogen; R₂ is hydrogen, COOH or CH₃; R₃ is hydrogen or COOH, R₂ being COOH or CH₃, when R₃ is hydrogen; and R₄ is hydrogen.

48. A compound according to claim 47 wherein R₂ is COOH and R₃ is hydrogen.

49. A compound according to claim 46 wherein R₁ and R₃ are H, R₂ is COOH or CH₃, and R₄ is a phenyl or n-oxypropyl group.

50. A compound according to claim 46 wherein R₁, R₃, and R₄ are H; and R₂ is tetrazolyl.

51. A compound according to claim 40,
wherein R_0 is $C_xH_y-SO_2-NH-$, x is 2-16 and y is 3 to 33.

52. A compound according to claim 51,
5 wherein R_0 is selected from the group consisting of $C_2H_5-SO_2-NH-$, $CH_2=CH-SO_2-NH-$, and $CH_3(CH_2)_{15}-SO_2-NH-$.

53. A compound according to claim 40,
wherein R_0 is NH_2- .

54. A compound according to claim 53,
10 wherein R is a 2'-carboxy-biphenyl group or 1 methoxy-2'-carboxy-biphenyl group.

55. A compound according to claim 39 wherein
 R_0 is SH and B is 3 aminomethyl-3'-carboxy-biphenyl.

56. A method of inhibiting p21ras
15 farnesyltransferase in a host in need of such inhibition which comprises administering to said host an effective amount of a compound according to claim 39.

57. A pharmaceutical composition comprising
20 a compound according to claim 39 and a pharmaceutically acceptable carrier therefor.

58. A method of inhibiting p21ras
25 farnesyltransferase in a host in need of such inhibition which comprises administering to said host an effective amount of a compound according to claim 41.

59. A pharmaceutical composition comprising a compound according to claim 41 and a pharmaceutically acceptable carrier therefor.

60. A peptidomimetic of formula:

5

 $C\beta X$

wherein

C represents cysteine;

X is an amino acid; and

10 β is a non-peptide aminoalkyl- or amino-substituted aliphatic or aromatic carboxylic acid or a heterocyclic monocarboxylic acid.

61. A peptidomimetic according to claim 60 wherein β is an aminobenzoic or aminoalkanoic acid.

15

62. A peptidomimetic of the formula:

 CA

wherein C is cysteine and Δ is an aryl or heterocyclic substituent which does not include a peptide amino acid but corresponds essentially in size with a tetrapeptide of the formula CA_1A_2X wherein C is cysteine and A_1 , A_2 and X are peptide amino acids.

20

63. A peptidomimetic according to claim 65 comprising cysteine joined directly to an aminomethyl-bicyclic arylcarboxylic acid.

25

64. A compound according to claim 63 wherein Δ comprises a biphenyl group.


65. A compound according to claim 64 wherein Δ is the radical of 3-aminomethyl-biphenyl-3-carboxylic acid.

66. A compound of the formula:

5



or



wherein

10

C is cysteine;

A_1 , A_2 and X are peptide amino acids;

Δ is an aryl or heterocyclic substituent which does not include a peptide amino acid;

15

β is a non-peptide aminoalkyl- or amino-substituted aliphatic or aromatic carboxylic acid or heterocarboxylic acid; and

20

Δ_1 is a phosphonate group joined to C through the cysteine S atom.

67. A compound according to claim 66 of the formula:




25 wherein Δ_1 , C and X have the meanings given in claim 56 and AMBA is an aminomethyl-benzoic acid.


68. A compound according to claim 67 wherein AMBA is 3-aminomethyl benzoic acid.

69. A compound of the formula:

30



or



wherein

C is cysteine; A₁, A₂ and X are peptide amino acids;

5 Δ is an aryl or heterocyclic substituent which does not include a peptide amino acid;

 β is a non-peptide aminoalkyl- or amino-substituted aliphatic or aromatic carboxylic acid or
10 heterocarboxylic acid; and

 β₁ is an aryl or aralkyl group joined to C through the cysteine S atom.



70. A compound according to claim 69 of the formula:

15 β₁ C-AMBA-X


wherein C, X and β₁ have the meanings given in claim 70 and AMBA is an aminomethyl benzoic acid.

71. A compound according to claim 70 wherein AMBA is 3-aminomethylbenzoic acid.

72. A compound of the formula:



or







73. A method of improving peptide or peptidomimetic uptake into cells which comprises utilizing the peptide in a functionalized form such that terminal amino, sulfhydryl and acid groups are functionalized by lipophilic or hydrophobic, esterase-sensitive moieties.

74. A method according to claim 73 for inhibiting tumor cell growth in a host which comprises administering to the host an effective amount of a peptide or peptidomimetic wherein one or more terminal amino, sulfhydryl and carboxylic acid groups have been functionalized with a lipophilic or hydrophobic esterase-sensitive moiety.

75. A method according to claim 74 wherein the terminal amino and sulfhydryl groups are functionalized by benzyloxy carbonyl groups and the carboxylic acid group is esterified.

76. A compound represented by one of formulas (A-L):




wherein R_1' is

- i) hydrogen;
- ii) lower alkyl;
- iii) alkenyl;
- 5 iv) alkoxy;
- v) thioalkoxy;
- vi) halo;
- vii) haloalkyl;
- viii) aryl- L_2 -, wherein L_2 is absent, $-CH_2-$, $-CH_2CH_2-$, $-CH(CH_3)-$, $-O-$, $-S(O)_q$ wherein q is 0, 1, or 2, $-N(R')$ - wherein R' is hydrogen or lower alkyl, or $-C(O)-$ and aryl is selected from the group consisting of phenyl, naphthyl, tetrahydronaphthyl, indanyl and indenyl and the
- 10 aryl group is unsubstituted or substituted; or
- ix) heterocyclic- L_3 - wherein L_3 is absent, $-CH_2-$, $-CH_2CH_2-$, $-CH(CH_3)-$, $-O-$, $-S(O)_q$ wherein q is 0, 1 or 2, $-N(R')$ - wherein R' is hydrogen or loweralkyl, or $-C(O)-$ and heterocyclic is a
- 15 monocyclic heterocyclic wherein the heterocyclic is unsubstituted or substituted with one, two, or three substituents independently selected from the group consisting of loweralkyl, hydroxy, hydroxyalkyl, halo, nitro, oxo ($=O$), amino, N-
- 20 protected amino, alkoxy, thioalkoxy and haloalkyl;
- 25

R_{1a} is hydrogen or lower alkyl;

R_2' is

i)



- 30 wherein R_{12a} is hydrogen, loweralkyl or $-C(O)O-R_{13}$, wherein R_{13} is hydrogen or a carboxy-protecting



group and R_{12b} is hydrogen or loweralkyl, with the proviso that R_{12a} and R_{12b} are not both hydrogen,

ii) $-C(O)NH-CH(R_{14})-C(O)OR_{15}$ wherein R_{14} is

- 5 a) loweralkyl,
b) cycloalkyl,
c) cycloalkylalkyl,
d) alkoxyalkyl,
e) thioalkoxyalkyl,
f) hydroxyalkyl,
- 10 g) aminoalkyl,
h) carboxyalkyl,
i) alkoxycarbonylalkyl,
j) arylalkyl or
k) alkylsulfonylalkyl and

15 R_{15} is hydrogen or a carboxy-protecting group or

iii)

 R_3' is

where

25 A represents O or 2H, and
R₀ represents SH, NH₂, or C_xH_y-SO₂-NH-, wherein
C_xH_y is a straight chain saturated or unsaturated

hydrocarbon, with x being between 1 and 20 and y between 3 and 41, inclusive; and

B stands for -NHR, where R is an aryl group.

and

- 5 L_1 is -NH-;
or pharmaceutically acceptable salts or prodrugs thereof.

1 / 22

FIG. 1A





FIG. 1B



2 / 22




FIG. 2A



FIG. 2B

3 / 22




FIG. 3

4 / 22




FIG. 4A




FIG. 4B SUBSTITUTE SHEET (RULE 26)

5 / 22




FIG. 5A




FIG. 5B

6 / 22




FIG. 6

FIG. 7A

7 / 22





FIG. 7B



SUBSTITUTE SHEET (RULE 26) Days

8 / 22

FIG. 8A





FIG. 8B



9 / 22

FIG. 9A





FIG. 9B



10 / 22

FIG. 10A





FIG. 10B



11 / 22




FIG. IIA








FIG. IIB SUBSTITUTE SHEET (RULE 26)

12 / 22



FIG. 12

4 : R¹=COOH R²=H5 : R¹=H R²=COOH8 : R¹=CH₃ R²=H

SUBSTITUTE SHEET (RULE 26)

13 / 22

FIG. 13



14 / 22

FIG. 14A





FIG. 14B



15 / 22




FIG. 15

16 / 22


Ras

FIG. 16A


Rap1A

FIG. 16B

17 / 22






FIG. 17

FTI-276: $R = O^-$ FTI-277: $R = OCH_3$ GGTI-286: $R = OCH_3$ GGTI-287: $R = O^-$

GGTI-297: red. 2-(1-Naphthyl) CABAL



SUBSTITUTE SHEET (RULE 26)

18 / 22




FIG. 18

19 / 22

K-Ras4B




FIG. 19

20 / 22




K-Ras4B**H-Ras**

FIG. 20


21 / 22

FIG. 21



22 / 22

FIG. 22



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/01559

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :A61K 38/06 US CL :514/18 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols), U.S. : 514/18, 19; 562/557 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS Online, APS		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Cell, Volume 62, issued 13 July 1990, Y. Reiss et al, "Inhibition of Purified p21 ^{ras} Farnesyl:Protein Transferase by Cys-AAX Tetrapeptides", pages 81-88, see entire document.	1-76
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* "A" "E" "L" "T" "I"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	"T" "X" "Y" "G" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
Date of the actual completion of the international search 31 MAY 1996		Date of mailing of the international search report 21 JUN 1996
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer DAVID LUKTON <i>David Lukton</i> Telephone No. (703) 308-0196